

1 The heat equation.

Recall that if we study the second order equation

$$au_{xx} + 2bu_{xy} + cu_{yy} + 2du_x + 2eu_y + ku = f(x, y), \quad u = u(x, y) \quad (1)$$

where a, \dots, k are all **constants** and $f(x, y)$ is a given function defined on some open set $\Omega \subseteq \mathbb{R}^2$, then we have the following classification result:

Theorem 1.1 (Refined canonical form.) *If the linear equation (1) is **elliptic**, then one can find a suitable linear change of variables (using eigenvalues, eigenvectors and scalings) and multiply the solution by some **exponential function** so that, eventually, the equation has the form*

$$v_{\xi\xi} + v_{\eta\eta} + cv = \phi(\xi, \eta), \quad v = v(\xi, \eta), \quad (2)$$

for some constant $c \in (-\infty, \infty)$ and some function $\phi(\xi, \eta)$. If the equation (1) is **hyperbolic**, the equation has the form

$$v_{\xi\xi} - v_{\eta\eta} + cv = \phi(\xi, \eta), \quad v = v(\xi, \eta), \quad (3)$$

for some constant $c \in (-\infty, \infty)$ and some function $\phi(\xi, \eta)$. If the equation (1) is **parabolic** and **nondegenerate**, the equation has the form

$$v_{\xi\xi} + cv_{\eta} = \phi(\xi, \eta), \quad v = v(\xi, \eta), \quad (4)$$

for some constant $c \neq 0$ and some function $\phi(\xi, \eta)$.

Remark 1.2 (Important.) *The constant c in the **elliptic case** can be $c > 0$, or $c = 0$, or $c < 0$. For $c > 0$, we can make it equal to 1 by doing the change of variables*

$$\tilde{\xi} = \sqrt{c}\xi, \quad \tilde{\eta} = \sqrt{c}\eta, \quad \tilde{v}(\tilde{\xi}, \tilde{\eta}) = v\left(\frac{\xi}{\sqrt{c}}, \frac{\eta}{\sqrt{c}}\right),$$

and for $c < 0$, we can make it equal to -1 by doing the change of variables

$$\tilde{\xi} = \sqrt{-c}\xi, \quad \tilde{\eta} = \sqrt{-c}\eta, \quad \tilde{v}(\tilde{\xi}, \tilde{\eta}) = v\left(\frac{\xi}{\sqrt{-c}}, \frac{\eta}{\sqrt{-c}}\right).$$

Thus in the **elliptic case**, we may simply assume $c = 1$, or 0 , or -1 . The constant c in the **hyperbolic case** can be $c > 0$, or $c = 0$, or $c < 0$. For $c < 0$, by switching the role of ξ and η , we may assume $c > 0$, or $c = 0$. Hence for the **hyperbolic case**, eventually, we can simply assume $c = 1$, or 0 . Finally, for the **parabolic case**, the constant $c \neq 0$ can be $c > 0$, or $c < 0$. So eventually we can simply assume $c = 1$, or -1 . However, since most parabolic equations come from physical phenomenon involving the behavior of some quantity $v(\xi, \eta)$ depending on space and time. So ξ will represent **space variable** and η will represent **time variable**. In that case a **model parabolic equation** looks like (assume $\phi(\xi, \eta) = 0$ for simplicity)

$$(1). v_t = v_{xx} \quad \text{or} \quad (2). v_t = -v_{xx}. \quad (5)$$

We call (1) the "**forward heat equation**" (or just **heat equation**) and (2) the "**backward heat equation**". Since in reality, time cannot go backwards, so in a parabolic equation, we always focus on the behavior of a solution $v(x, t)$ **as time goes forwards**, i.e., as t is **increasing**. One can

use simple examples to see that, **as time goes forwards**, the heat equation (1) will make solution better, while the backward heat equation (2) will make solution worse (look at $e^{-t} \sin x$ and $e^t \sin x$ respectively). Thus, as time goes forwards, equation (1) is well-posed, while (2) is ill-posed. In this course, we will focus only on (1) (on the other hand, **as time goes backwards**, (1) will make solution worse and (2) will make solution better...).

Definition 1.3 Let $v = v(\xi, \eta)$. The equations $v_{\xi\xi} + v_{\eta\eta} = 0$, $v_{\xi\xi} - v_{\eta\eta} = 0$, $v_{\xi\xi} - v_{\eta} = 0$, are called **Laplace equation (elliptic equation)**, **wave equation (hyperbolic equation)**, and **heat equation (nondegenerate parabolic equation)**, respectively.

By Theorem 1.1 and Remark 1.2, we study the following **one-dimensional heat equation (nondegenerate parabolic equation)** (we focus on equation (1) in (5)):

$$u_t = u_{xx}, \quad u = u(x, t), \quad x \in \mathbb{R}. \quad (6)$$

For higher dimensional heat equation, it has the form

$$u_t = \Delta u, \quad u = u(x_1, \dots, x_n, t), \quad (x_1, \dots, x_n) \in \mathbb{R}^n,$$

where

$$\Delta = \frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_n^2} \text{ is the Laplace operator in } \mathbb{R}^n.$$

We will focus only on the one-dimensional case, i.e. $n = 1$, in this course.

2 Physical motivation for the heat equation.

We will give a brief explanation why the equation $u_t = \Delta u$ is called the **heat equation**. The reason is that it describes the behavior of the **temperature function** in the **heat flow phenomenon**. We look at the case $n = 3$ and let $\Omega \subset \mathbb{R}^3$ be a bounded "**heated domain**". At any time $t \in (0, \infty)$, let $u(x, y, z, t)$ be the temperature at the point $\mathbf{x} = (x, y, z) \in \Omega$. The **total heat** inside the domain Ω at time $t \in (0, \infty)$ is given by

$$H(t) = \int_{\Omega} u(\mathbf{x}, t) d\mathbf{x}, \quad t \in (0, \infty). \quad (7)$$

The change of total heat inside Ω is given by (here we assume we can differentiate under the integral sign, which is actually so in most situations)

$$\frac{dH}{dt}(t) = \int_{\Omega} \frac{\partial u}{\partial t}(\mathbf{x}, t) d\mathbf{x}, \quad t \in (0, \infty). \quad (8)$$

On the other hand, by physical experiment, the French mathematician J. Fourier discovered that the heat will flow from hot to cold regions in a way that is **proportional to the gradient of the temperature everywhere**, i.e., proportional to the quantity

$$\nabla u = \left(\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial z} \right), \quad (9)$$

with certain proportion constant κ (heat conductivity).

Moreover, due to the **conservation law of energy**, if there is a change in the total heat inside Ω , it must be due to the heat flowing out or flowing into Ω through **the boundary** $\partial\Omega$ (which is a surface in \mathbb{R}^3). By **conservation law** and **Fourier's law**, we also have the identity

$$\frac{dH}{dt}(t) = \int_{\partial\Omega} \kappa (\nabla u(\mathbf{x}, t) \cdot \mathbf{N}(\mathbf{x})) dS \text{ (surface integral in } \mathbb{R}^3), \quad t \in (0, \infty), \quad (10)$$

where $\mathbf{N}(\mathbf{x})$ is the **unit outward normal** of $\partial\Omega$ at $\mathbf{x} \in \partial\Omega$ and $\nabla u(\mathbf{x}, t) \cdot \mathbf{N}(\mathbf{x})$ is the inner product in \mathbb{R}^3 between the two vectors $\nabla u(\mathbf{x}, t)$ and $\mathbf{N}(\mathbf{x})$. By (8) and (10) and the classical **divergence theorem**, we have the identity

$$\int_{\Omega} \frac{\partial u}{\partial t}(\mathbf{x}, t) d\mathbf{x} = \int_{\partial\Omega} \kappa (\nabla u(\mathbf{x}, t) \cdot \mathbf{N}(\mathbf{x})) dS = \int_{\Omega} \kappa \operatorname{div}(\nabla u(\mathbf{x}, t)) d\mathbf{x} = \int_{\Omega} \kappa \Delta u(\mathbf{x}, t) d\mathbf{x}$$

and so we conclude the **integral identity** on Ω :

$$\int_{\Omega} \left[\frac{\partial u}{\partial t}(\mathbf{x}, t) - \kappa \Delta u(\mathbf{x}, t) \right] d\mathbf{x} = 0, \quad \forall t \in (0, \infty). \quad (11)$$

Finally, we note that the analysis leading to the identity (11) is independent of the domain Ω , i.e. on any subdomain $\tilde{\Omega} \subset \Omega \subset \mathbb{R}^3$, as long as the heat flow phenomenon obeys the **conservation law** and **Fourier's law** in $\tilde{\Omega}$, we always have the identity

$$\int_{\tilde{\Omega}} \left[\frac{\partial u}{\partial t}(\mathbf{x}, t) - \kappa \Delta u(\mathbf{x}, t) \right] d\mathbf{x} = 0, \quad \forall t \in (0, \infty). \quad (12)$$

Since the domain $\tilde{\Omega} \subset \Omega \subset \mathbb{R}^3$ in (12) is arbitrary, we must have

$$\frac{\partial u}{\partial t}(\mathbf{x}, t) = \kappa \Delta u(\mathbf{x}, t), \quad \forall (\mathbf{x}, t) \in \Omega \times (0, \infty), \quad (13)$$

which is the heat equation if we do suitable scaling to make $\kappa = 1$.

3 The 1-dimensional heat equation.

Most of the time (but not always) we will focus only on the 1-dimensional heat equation in this course ("1-dimensional" means space dimension n is 1). Unlike the 1-dimensional wave equation $u_{tt} = u_{xx}$, the heat equation $u_t = u_{xx}$ is much more difficult to solve. It is not difficult to guess some special solutions of $u_t = u_{xx}$, like

$$u(x, t) = x, \quad 2t + x^2, \quad x^3 + 6xt, \quad e^{t+x}, \quad e^{t-x}, \quad e^{-t} \cos x, \quad e^{-t} \sin x, \quad e^t \cosh x, \quad e^t \sinh x, \quad (14)$$

, etc. (note that $e^t \cosh x$ and $e^t \sinh x$ are linear combinations of e^{t+x} and e^{t-x}). All of the above solutions are defined on $\mathbb{R} \times \mathbb{R}$. One can check that the only **space-time separable solutions** of the heat equation are "**essentially**" of the form

$$u(x, t) = 1, \quad x, \quad e^{t+x}, \quad e^{t-x}, \quad e^{-t} \cos x, \quad e^{-t} \sin x \quad (15)$$

and no others. Also note that the solutions $u(x, t) = x, 2t + x^2, x^3 + 6xt$ are **polynomial solutions** with $u(x, 0) = x, x^2, x^3$. There is a formula for a polynomial solution with $u(x, 0) = x^n$ for any $n \in \mathbb{N}$. We will discuss this later on.

There are several major differences between the wave equation and the heat equation:

1. There is smoothing effect for heat equation, but not so in wave equation. We will discuss this later on.
2. For wave equation, if $u(x, t)$ is a solution, so is the function $u(x, -t)$, but for the heat equation, if $u(x, t)$ is a solution, the function $u(x, -t)$ is, in general, no longer a solution. Thus for the heat equation $u_t = u_{xx}$, one cannot reverse the direction of time.
3. (**Scaling property**.) If $u(x, t)$ is a solution of the heat equation, so is the function $\tilde{u}(x, t) = u(\lambda x, \lambda^2 t)$ for any constant $\lambda \neq 0$ (for the wave equation, if $u(x, t)$ is a solution, so is the function $\tilde{u}(x, t) = u(\lambda x, \lambda t)$ for any constant $\lambda \neq 0$).

Example 3.1 (Interesting solutions.) We have the following interesting solutions of $u_t = u_{xx}$. They are all defined on $\mathbb{R} \times (-\infty, \infty)$.

$$u(x, t) = e^{-t} \cos x, \quad e^{-t} \sin x \quad (\text{space-periodic solutions, } u(x + 2\pi, t) = u(x, t))$$

and

$$u(x, t) = e^{\pm \frac{x}{\sqrt{2}}} \cos\left(t \pm \frac{x}{\sqrt{2}}\right), \quad e^{\pm \frac{x}{\sqrt{2}}} \sin\left(t \pm \frac{x}{\sqrt{2}}\right) \quad (\text{time-periodic solutions, } u(x, t + 2\pi) = u(x, t))$$

and

$$u(x, t) = \begin{cases} x^2 + 2t, & x^3 + 6xt, \\ x^4 + 12x^2t + 12t^2, & x^5 + 20x^3t + 60xt^2, \dots \end{cases} \quad (\text{polynomial solutions}),$$

where we note the important property that t is like x^2 (see Remark 3.7 below), and

$$u(x, t) = e^{t+x}, \quad e^{t-x} \quad (\text{traveling wave solutions}).$$

Note that a function $u(x, t)$ of the form $u(x, t) = h(x - \lambda t)$ for some constant $\lambda \in \mathbb{R}$ is usually called a **traveling wave function**. To understand this terminology, you can plot the graphs of $u(x, 0)$, $u(x, 1)$, $u(x, 2)$, $u(x, 3)$, ..., and see that the graph of $u(x, 0)$ is moving along the x -direction as time goes on.

3.1 Polynomial solutions of the 1-dimensional heat equation.

If we do not impose any "side condition" on the heat equation $u_t = u_{xx}$, then on \mathbb{R}^2 it has infinitely many solutions. Recall that for the Laplace equation on \mathbb{R}^2 , we have a family of polynomial solutions known as "harmonic polynomials". They are $1, x, y, xy, x^2 - y^2, \dots$, and they are all defined on \mathbb{R}^2 . In terms of polar coordinates (r, θ) in the plane they have the forms $r^n \cos n\theta, r^n \sin n\theta$ for $n \in \mathbb{N} \cup \{0\}$. These solutions are important because we can use them to construct the **Poisson Integral Formula** on the disc.

For the heat equation $u_t(x, t) = u_{xx}(x, t)$ on $(x, t) \in \mathbb{R}^2$, there are also "heat polynomials" defined on the whole space $(x, t) \in \mathbb{R}^2$. In below, we show you how to derive them.

Consider the 1-dimensional heat equation $u_t(x, t) - u_{xx}(x, t) = 0$ with **initial data** (data at $t = 0$)

$$u(x, 0) = p_0(x), \quad x \in (-\infty, \infty), \tag{16}$$

where $p_0(x)$ is a **polynomial** defined on $x \in (-\infty, \infty)$ with degree $n \in \mathbb{N} \cup \{0\}$. We try to look for a **space-time polynomial solution** $u(x, t)$ of the heat equation of the form

$$u(x, t) = p_0(x) + p_1(x)t + p_2(x)t^2 + p_3(x)t^3 + \dots,$$

where each $p_i(x)$ is also a polynomial in $x \in (-\infty, \infty)$.

We compute

$$u_t(x, t) = p_1(x) + 2p_2(x)t + 3p_3(x)t^2 + \dots$$

and

$$u_{xx}(x, t) = p_0''(x) + p_1''(x)t + p_2''(x)t^2 + p_3''(x)t^3 + \dots,$$

and by comparing the coefficient functions (because we want $u_t(x, t) = u_{xx}(x, t)$), we require

$$\left\{ \begin{array}{l} p_1(x) = p_0''(x), \\ p_2(x) = \frac{1}{2}p_1''(x) = \frac{1}{2}p_0''''(x), \\ p_3(x) = \frac{1}{3}p_2''(x) = \frac{1}{3!}p_0^{(6)}(x), \\ \dots \\ p_k(x) = \frac{1}{k}p_{k-1}''(x) = \frac{1}{k!}p_0^{(2k)}(x), \\ \dots \end{array} \right. \quad (17)$$

Since $p_0(x)$ is a polynomial with finite degree $n \in \mathbb{N}$, the above process will stop at some k (i.e. $p_0^{(2k)}(x)$ will become 0 for some $k \in \mathbb{N}$). Moreover, we see that all of the other polynomials $p_1(x)$, $p_2(x)$, $p_3(x)$, ..., can be **uniquely determined** by $p_0(x)$, which is the **initial condition** of the heat equation. Therefore, if the polynomial $p_0(x)$ is given in advance, we can find a **unique polynomial solution** of the heat equation $u_t = u_{xx}$ defined on $(x, t) \in \mathbb{R}^2$ satisfying (16).

We look at some simple examples.

Example 3.2 Take $p_0(x) = x$. Then $p_1(x) = p_0''(x) = 0$ and so on. The function $u(x, t) = x$ is a polynomial solution of the heat equation.

Example 3.3 Take $p_0(x) = x^2$. Then $p_1(x) = p_0''(x) = 2$ and $p_2(x) = 0$ and so on. The function

$$u(x, t) = p_0(x) + p_1(x)t = x^2 + 2t \quad (18)$$

is a polynomial solution of the heat equation.

Example 3.4 Take $p_0(x) = x^3$. Then $p_1(x) = p_0''(x) = 6x$ and $p_2(x) = 0$ and so on. The function

$$u(x, t) = p_0(x) + p_1(x)t = x^3 + 6xt \quad (19)$$

is a polynomial solution of the heat equation.

Example 3.5 Take $p_0(x) = x^4$. Then $p_1(x) = p_0''(x) = 12x^2$ and $p_2(x) = 12$ and $p_3(x) = 0$ and so on. The function

$$u(x, t) = p_0(x) + p_1(x)t + p_2(x)t^2 = x^4 + 12x^2t + 12t^2 \quad (20)$$

is a polynomial solution of the heat equation.

Example 3.6 Take $p_0(x) = x^5$. Then $p_1(x) = p_0''(x) = 20x^3$ and $p_2(x) = 60x$ and $p_3(x) = 0$ and so on. The function

$$u(x, t) = p_0(x) + p_1(x)t + p_2(x)t^2 = x^5 + 20x^3t + 60xt^2 \quad (21)$$

is a polynomial solution of the heat equation.

Remark 3.7 In all of the above examples, note that t is like x^2 (so that each term has the same degree !!). Therefore, in the solution

$$u(x, t) = x^5 + 20x^3t + 60xt^2,$$

we see that each term has "degree 5".

Remark 3.8 (You will understand this remark later on.) If we use the **representation formula** (you will see it later on)

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} p_0(y) dy, \quad t > 0, \quad (22)$$

we will get the **same** answer on the domain $(x, t) \in (-\infty, \infty) \times (0, \infty)$. Note that the integral (22) converges for any polynomial $p_0(y)$. Moreover, differentiation can move into the integral sign.

3.2 Finding the fundamental solution of the heat equation with the help of polynomial solutions.

Until now, we have found lots of **polynomial solutions** of the heat equation $u_t = u_{xx}$ on $(x, t) \in \mathbb{R}^2$, namely

$$x^2 + 2t, \quad x^3 + 6xt, \quad x^4 + 12x^2t + 12t^2, \quad x^5 + 20x^3t + 60xt^2, \quad \dots, \text{ etc.} \quad (23)$$

Restricted onto the domain $\mathbb{R} \times (0, \infty)$, each of the polynomial solution can be expressed as the form

$$u(x, t) = g(t) h\left(\frac{x^2}{t}\right), \quad (x, t) \in \mathbb{R} \times (0, \infty)$$

for some functions $g(t)$, $h(\theta)$ defined on $t \in (0, \infty)$, $\theta \in [0, \infty)$. For example, we can express

$$x^2 + 2t = g(t) h\left(\frac{x^2}{t}\right), \quad \text{where } g(t) = t, \quad h(\theta) = \theta + 2$$

and

$$\begin{aligned} x^3 + 6xt &= t^{3/2} \left(\left(\frac{x}{\sqrt{t}}\right)^3 + 6 \left(\frac{x}{\sqrt{t}}\right) \right) \\ &= g(t) h\left(\frac{x^2}{t}\right), \quad \text{where } g(t) = t^{3/2}, \quad h(\theta) = (\sqrt{\theta})^3 + 6\sqrt{\theta} \end{aligned}$$

and

$$\begin{aligned} x^4 + 12x^2t + 12t^2 &= t^2 \left(\left(\frac{x^2}{t}\right)^2 + 12 \left(\frac{x^2}{t}\right) + 12 \right) \\ &= g(t) h\left(\frac{x^2}{t}\right), \quad \text{where } g(t) = t^2, \quad h(\theta) = \theta^2 + 12\theta + 12. \end{aligned}$$

Therefore, we can plug the general form $u(x, t) = g(t) h\left(\frac{x^2}{t}\right)$ into the heat equation $u_t = u_{xx}$ and see if we can find new interesting solutions. Compute

$$u_t(x, t) = g'(t) h\left(\frac{x^2}{t}\right) - g(t) h'\left(\frac{x^2}{t}\right) \frac{x^2}{t^2}, \quad u_x(x, t) = g(t) h'\left(\frac{x^2}{t}\right) \frac{2x}{t}$$

and

$$u_{xx}(x, t) = g(t) h''\left(\frac{x^2}{t}\right) \frac{4x^2}{t^2} + g(t) h'\left(\frac{x^2}{t}\right) \frac{2}{t}.$$

We hope to have the identity

$$\underbrace{g'(t) h\left(\frac{x^2}{t}\right)} - \underbrace{g(t) h'\left(\frac{x^2}{t}\right) \frac{x^2}{t^2}} = \underbrace{g(t) h''\left(\frac{x^2}{t}\right) \frac{4x^2}{t^2}} + \underbrace{g(t) h'\left(\frac{x^2}{t}\right) \frac{2}{t}}, \quad \forall (x, t) \in \mathbb{R} \times (0, \infty), \quad (24)$$

which is **possible** if we require

$$\begin{cases} -h'(\theta) = 4h''(\theta), & \theta \in [0, \infty) \\ g'(t) h(\theta) = \left(g(t) \frac{2}{t}\right) h'(\theta), & \theta \in [0, \infty), \quad t \in (0, \infty). \end{cases} \quad (25)$$

Solving the first equation, we get the general solution $h(\theta) = A + Be^{-\frac{\theta}{4}}$ for arbitrary constants A , B and we choose $A = 0$, $B = 1$ and plug $h(\theta) = e^{-\frac{\theta}{4}}$ into the second equation to get the equation for g :

$$g'(t) = -\frac{1}{2t}g(t), \quad (26)$$

which gives the general solution $g(t) = \frac{C}{\sqrt{t}}$ for arbitrary constant C . Therefore, we see that

$$u(x, t) = g(t) h\left(\frac{x^2}{t}\right) = \frac{1}{\sqrt{t}} e^{-\frac{x^2}{4t}}, \quad (x, t) \in \mathbb{R} \times (0, \infty) \quad (27)$$

is a **new solution** of the heat equation on $\mathbb{R} \times (0, \infty)$. Note that this solution is different from any solution you encountered before. \square

Remark 3.9 If $g(t)$ and $h(\theta)$ are from a polynomial solution $u(x, t)$, then they will satisfy (24) too.

Remark 3.10 (Important.) If we use the fact: if $u_j(x, t)$ is a solution for the one-dimensional heat equation $u_t = u_{xx}$ on $\mathbb{R} \times (0, \infty)$, then the function

$$u(\mathbf{x}, t) = u_1(x_1, t) \cdots u_n(x_n, t), \quad \mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n \quad (28)$$

is a solution of the heat equation $u_t = \Delta u$ on $\mathbb{R}^n \times (0, \infty)$. With this, by (27), we will obtain the solution

$$u(\mathbf{x}, t) = \frac{1}{\sqrt{t}} e^{-\frac{x_1^2}{4t}} \cdots \frac{1}{\sqrt{t}} e^{-\frac{x_n^2}{4t}} = \frac{1}{t^{n/2}} e^{-\frac{|\mathbf{x}|^2}{4t}}, \quad \mathbf{x} \in \mathbb{R}^n, \quad t > 0 \quad (29)$$

of the heat equation $u_t = \Delta u$ on $\mathbb{R}^n \times (0, \infty)$.

By (29), we now define the following (for **normalization** purpose, we divide the solution in (29) by the constant $(4\pi)^{n/2}$; see Lemma 3.17):

Definition 3.11 The function

$$\Phi(\mathbf{x}, t) = \begin{cases} \frac{1}{(4\pi t)^{n/2}} e^{-\frac{|\mathbf{x}|^2}{4t}}, & \mathbf{x} \in \mathbb{R}^n, \quad t > 0 \\ 0, & \mathbf{x} \in \mathbb{R}^n, \quad t \leq 0. \end{cases} \quad (30)$$

is called the **fundamental solution** of the heat equation. For each fixed time t , it is **radial** in $\mathbf{x} \in \mathbb{R}^n$. Moreover it satisfies the heat equation $\partial_t \Phi = \Delta \Phi$ in $\mathbb{R}^{n+1} \setminus \{(0, 0)\}$ and is **invariant** under the **space-time scaling** $\Phi(\mathbf{x}, t) \rightarrow \lambda^n \Phi(\lambda \mathbf{x}, \lambda^2 t)$, i.e. we have

$$\lambda^n \Phi(\lambda \mathbf{x}, \lambda^2 t) = \Phi(\mathbf{x}, t), \quad \forall \lambda > 0, \quad \forall (\mathbf{x}, t) \in \mathbb{R}^{n+1}. \quad (31)$$

Remark 3.12 (1). The **only singularity** of Φ is at the point $(0, 0)$, i.e. $\Phi(\mathbf{x}, t) \in C^\infty(\mathbb{R}^{n+1} \setminus \{(0, 0)\})$ and it is **not continuous** at $(0, 0)$. To understand the property $\Phi(\mathbf{x}, t) \in C^\infty(\mathbb{R}^{n+1} \setminus \{(0, 0)\})$ you need to know the fact that for each fixed $\mathbf{x}_0 \neq 0 \in \mathbb{R}^n$, the function

$$\psi(t) = \begin{cases} \frac{1}{(4\pi t)^{n/2}} e^{-\frac{|\mathbf{x}_0|^2}{4t}}, & t > 0 \\ 0, & t \leq 0 \end{cases} \quad (32)$$

is a C^∞ function of $t \in (-\infty, \infty)$. On the other hand, for $\mathbf{x}_0 = 0$, $\psi(t)$ becomes

$$\psi(t) = \begin{cases} \frac{1}{(4\pi t)^{n/2}}, & t > 0 \\ 0, & t \leq 0, \end{cases}$$

with $\lim_{t \rightarrow 0^+} \psi(t) = +\infty$ and so it is not continuous at $t = 0$. (2). $\Phi(\mathbf{x}, t)$ satisfies the heat equation $\partial_t \Phi = \Delta \Phi$ in $\mathbb{R}^{n+1} \setminus \{(0, 0)\}$. There is an easy way to check this on $\mathbb{R}^n \times (0, \infty)$. Let

$v = \ln \Phi$. Then Φ satisfies the heat equation $\partial_t \Phi = \Delta \Phi$ if and only if $v = \ln \Phi$ satisfies the equation $\partial_t v = \Delta v + |\nabla v|^2$ (this is an exercise for you to check). Therefore we check the later equation. We have

$$v = \ln \Phi = -\frac{n}{2} \ln(4\pi t) - \frac{|\mathbf{x}|^2}{4t}, \quad t > 0.$$

and then

$$\frac{\partial v}{\partial t} = -\frac{n}{2t} + \frac{|\mathbf{x}|^2}{4t^2}.$$

Also

$$\Delta v = -\frac{n}{2t}, \quad |\nabla v|^2 = \frac{|\mathbf{x}|^2}{4t^2}.$$

Hence we have $\partial_t v = \Delta v + |\nabla v|^2$. (3). Exercise: check that we have $\partial_t \Phi(\mathbf{x}, 0) = \Delta \Phi(\mathbf{x}, 0)$ for all $\mathbf{x} \in \mathbb{R}^n \setminus \{0\}$, $\mathbf{x} \neq 0$.

3.3 Basic properties of the fundamental solution.

In order to study the **initial value problem** for the heat equation (see (59) below) and to derive its solution formula, we need to discuss several important properties for the fundamental solution $\Phi(\mathbf{x}, t)$ given in (30). One can use this fundamental solution to give a **representation formula (solution formula)** for the solution of (59) (this is similar to the **Poisson Integral Formula** for Laplace equation on the disc).

As a comparison, recall that for the Laplace equation $\Delta u(\mathbf{x}) = 0$ in \mathbb{R}^n there is a **radial solution** (with a **singularity** at the origin of \mathbb{R}^n , i.e. $\mathbf{x} = 0$) of the form

$$u(\mathbf{x}) = \begin{cases} A|\mathbf{x}|^{2-n} + B, & n > 2, \text{ where } \mathbf{x} \in \mathbb{R}^n \setminus \{0\} \\ A \log |\mathbf{x}| + B, & n = 2, \text{ where } \mathbf{x} \in \mathbb{R}^2 \setminus \{0\}, \end{cases}$$

where A, B are arbitrary constants. It plays an important role in the theory of Laplace equation.

For the heat equation $u_t = \Delta u$, the fundamental solution $\Phi(\mathbf{x}, t)$ given in (30) is also a **radial solution (radial in space \mathbb{R}^n , not in space-time \mathbb{R}^{n+1})**, which, similar to the elliptic case, has a **singularity at the origin** of \mathbb{R}^{n+1} , i.e. at $(x, t) = (0, 0)$.

In the following, we will discuss several properties of the fundamental solution $\Phi(\mathbf{x}, t)$ for the case $n = 1$. These properties are all valid for general $n > 1$, but for simplicity of proof, here we focus only on the case $n = 1$.

Lemma 3.13 *Let*

$$\Phi(x, t) = \begin{cases} \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}}, & x \in \mathbb{R}, \quad t > 0 \\ 0, & x \in \mathbb{R}, \quad t \leq 0. \end{cases} \quad (33)$$

Then $\Phi(x, t) \in C^\infty(\mathbb{R}^2 \setminus \{(0, 0)\})$ and it satisfies the heat equation $\partial_t u = \Delta u$ in $\mathbb{R}^2 \setminus \{(0, 0)\}$.

Proof. Since

$$\Phi(0, t) = \begin{cases} \frac{1}{\sqrt{4\pi t}}, & t > 0 \\ 0, & t \leq 0, \end{cases}$$

we see that $\Phi(x, t)$ is *not* continuous at $(0, 0)$. Moreover, we have

$$\lim_{t \rightarrow 0^+} \Phi(0, t) = \lim_{t \rightarrow 0^+} \frac{1}{\sqrt{4\pi t}} = \infty.$$

To check that $\Phi(x, t) \in C^\infty(\mathbb{R}^2 \setminus \{(0, 0)\})$, it suffices to look at the behavior of $\Phi(x, t)$ on the set $S = \{(x, 0) \in \mathbb{R}^2 : x \neq 0\}$. By the limit

$$\lim_{t \rightarrow 0^+} \left(\frac{1}{t^\alpha} e^{-\frac{\beta}{t}} \right) = 0, \quad \forall \text{ const. } \alpha, \beta > 0,$$

one can check that $\Phi(x, t)$ is C^∞ at any point of S . Computing

$$\begin{aligned} \frac{\partial}{\partial x} \left(\frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} \right) &= \frac{1}{\sqrt{4\pi t}} \left(-\frac{x}{2t} \right) e^{-\frac{x^2}{4t}}, \quad t > 0 \\ \frac{\partial^2}{\partial x^2} \left(\frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} \right) &= \frac{1}{\sqrt{4\pi t}} \left(-\frac{1}{2t} + \frac{x^2}{4t^2} \right) e^{-\frac{x^2}{4t}}, \quad t > 0 \\ \frac{\partial}{\partial t} \left(\frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} \right) &= \frac{1}{\sqrt{4\pi}} \left(-\frac{1}{2} t^{-3/2} \right) e^{-\frac{x^2}{4t}} + \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} \frac{x^2}{4t^2}, \quad t > 0 \end{aligned} \quad (34)$$

we see that $\Phi(x, t)$ satisfies the heat equation on $\mathbb{R} \times (0, \infty)$. Clearly it also satisfies the heat equation on $\mathbb{R} \times (-\infty, 0)$. At any point $(x_0, 0) \in S$, $x_0 \neq 0$, we have $\Phi_{xx}(x_0, 0) = 0$. Also note that

$$\lim_{t \rightarrow 0^-} \frac{\Phi(x_0, t) - \Phi(x_0, 0)}{t} = 0 \quad (x_0 \neq 0)$$

and

$$\lim_{t \rightarrow 0^+} \frac{\Phi(x_0, t) - \Phi(x_0, 0)}{t} = \lim_{t \rightarrow 0^+} \left(\frac{1}{t} \frac{1}{\sqrt{4\pi t}} e^{-\frac{x_0^2}{4t}} \right) = 0 \quad (x_0 \neq 0, x_0^2 > 0),$$

and so we have $\Phi_t(x_0, 0) = 0$. The proof is done. \square

Lemma 3.14 *For any fixed $\varepsilon > 0$, we have*

$$\lim_{t \rightarrow 0^+} \Phi(x, t) = 0 \quad \text{uniformly in the region } \{x \in \mathbb{R} : |x| \geq \varepsilon\}. \quad (35)$$

Also

$$\lim_{|x| \rightarrow \infty} \Phi(x, t) = 0 \quad \text{uniformly in the region } t \in (-\infty, \infty). \quad (36)$$

Remark 3.15 *We also have*

$$\lim_{t \rightarrow \infty} \Phi(x, t) = 0 \quad \text{uniformly in } x \in (-\infty, \infty). \quad (37)$$

This is easy due to

$$|\Phi(x, t)| = \left| \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} \right| \leq \frac{1}{\sqrt{4\pi t}} \quad \text{for all } x \in (-\infty, \infty), \quad t > 0. \quad (38)$$

Remark 3.16 *Draw a picture for $\Phi(x, t)$ with $t \rightarrow 0^+$.*

Proof. For (35), we have for $t > 0$ the inequality

$$0 < \Phi(x, t) = \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} \leq \frac{1}{\sqrt{4\pi t}} e^{-\frac{\varepsilon^2}{4t}}, \quad \forall |x| \geq \varepsilon$$

and the conclusion follows. For (36), it suffices to focus on $t \in (0, \infty)$ since $\Phi(x, t) \equiv 0$ for all $x \in \mathbb{R}$, $t \leq 0$. For fixed $x \in \mathbb{R}$, $x \neq 0$, the maximum value of the positive function

$$\frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}}, \quad t \in (0, \infty)$$

over $t \in (0, \infty)$, is attained at the point $t = x^2/2$ with maximum value equal to

$$\frac{1}{|x| \sqrt{2\pi}} e^{-\frac{1}{2}}. \quad (39)$$

This is due to the identity

$$\frac{\partial}{\partial t} \left(\frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} \right) = \frac{1}{\sqrt{4\pi t}} \frac{1}{2t} \left(\frac{x^2}{2t} - 1 \right) e^{-\frac{x^2}{4t}}, \quad x \in \mathbb{R}, \quad t \in (0, \infty).$$

The result follows. \square

Lemma 3.17 *For each fixed $t > 0$, we have*

$$\int_{-\infty}^{\infty} \Phi(x, t) dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} dx = 1, \quad t > 0. \quad (40)$$

Moreover, the convergence of the integral is **uniform** with respect to $t \in (0, T)$ for any fixed $T > 0$ (but not uniform with respect to $t \in (0, \infty)$).

Remark 3.18 *Draw a picture for $\Phi(x, t)$ (for small $t > 0$ and for large $t > 0$) and show the property $\int_{-\infty}^{\infty} \Phi(x, t) dx = 1$ for all $t > 0$.*

Remark 3.19 (Helpful interpretation ...) *For fixed $t > 0$, if we let*

$$F_N(t) = \int_{-N}^N \Phi(x, t) dx, \quad t \in (0, T), \quad N \in \mathbb{N},$$

then the convergence of the integral is **uniform** with respect to $t \in (0, T)$ can be interpreted as

$$\lim_{N \rightarrow \infty} F_N(t) = 1 \quad \text{uniformly in } t \in (0, T).$$

Remark 3.20 *Similarly, we have*

$$\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dx = 1, \quad \forall y \in \mathbb{R}, \quad t > 0 \quad (41)$$

and

$$\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy = 1, \quad \forall x \in \mathbb{R}, \quad t > 0. \quad (42)$$

Proof. We first recall the following improper integral identity from calculus:

$$\int_{-\infty}^{\infty} e^{-s^2} ds = \sqrt{\pi}. \quad (43)$$

By a change of variables (let $s = \alpha y + \beta$), we have

$$\int_{-\infty}^{\infty} e^{-(\alpha y + \beta)^2} dy = \frac{\sqrt{\pi}}{\alpha}, \quad \forall \beta \in \mathbb{R}, \quad \alpha > 0. \quad (44)$$

Letting $x = \sqrt{4t}s$, we obtain

$$\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-s^2} \sqrt{4t} ds = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-s^2} ds = 1.$$

Next, let $T > 0$ be a fixed time. For any $\varepsilon > 0$, then there exists a large $M > 0$ (M depends only on ε and T) such that **for all** $t \in (0, T)$ we have the estimate (again, let $x = \sqrt{4t}s$)

$$0 < \int_M^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} dx = \frac{1}{\sqrt{\pi}} \int_{\frac{M}{\sqrt{4t}}}^{\infty} e^{-s^2} ds < \frac{1}{\sqrt{\pi}} \int_{\frac{M}{\sqrt{4T}}}^{\infty} e^{-s^2} ds < \varepsilon, \quad \forall t \in (0, T).$$

The same result holds for the integral $\int_{-\infty}^{-M} \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}} dx$. Therefore, the convergence of the integral is uniform with respect to $t \in (0, T)$ for any fixed $T > 0$. \square

Remark 3.21 By the integral identity

$$\int_{\mathbb{R}^n} e^{-|\mathbf{x}|^2} d\mathbf{x} = (\sqrt{\pi})^n,$$

one can also obtain the identity

$$\int_{\mathbb{R}^n} \frac{1}{(4\pi t)^{n/2}} e^{-\frac{|\mathbf{x}|^2}{4t}} d\mathbf{x} = 1 \quad (45)$$

for each $t > 0$.

Lemma 3.22 For **fixed** $\delta > 0$, we have

$$\lim_{t \rightarrow 0^+} \int_{|y-x| > \delta} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy = 0 \quad \text{uniformly in } x \in \mathbb{R}, \quad (46)$$

which means that the values of the fundamental solution $\Phi(x-y, t)$ (view it as a function of y with parameter x) concentrate around x as $t \rightarrow 0^+$.

Remark 3.23 For fixed $\delta > 0$, the quantity

$$\int_{|y-x| > \delta} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy$$

is a function of $(x, t) \in \mathbb{R} \times (0, \infty)$ (denote it as $F(x, t)$). The above lemma says that $\lim_{t \rightarrow 0^+} F(x, t) = 0$ **uniformly** in $x \in \mathbb{R}$.

Proof. Let $y = x + \sqrt{4t}s$. Then

$$\lim_{t \rightarrow 0^+} \int_{|y-x| > \delta} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy = \lim_{t \rightarrow 0^+} \frac{1}{\sqrt{\pi}} \int_{|s| > \delta/\sqrt{4t}} e^{-s^2} ds = 0. \quad (47)$$

Note that the right hand side of (47) does not depend on $x \in \mathbb{R}$. Hence we have convergence to zero uniformly in $x \in \mathbb{R}$. The proof is done. \square

The following lemma is crucial in solving the initial value problem (59) below.

Lemma 3.24 Let $\phi(x)$ be a **bounded** function defined on $(-\infty, \infty)$ and is **continuous** at $x = x_0$. Then we have

$$\lim_{(x,t) \rightarrow (x_0, 0^+)} \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy = \phi(x_0). \quad (48)$$

In particular, we also have

$$\lim_{t \rightarrow 0^+} \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x_0-y)^2}{4t}} \phi(y) dy = \phi(x_0). \quad (49)$$

Remark 3.25 The above two limits have different meaning. In the first limit, $(x, t) \rightarrow (x_0, 0^+)$ means that $(x, t) \in \mathbb{R} \times (0, \infty)$ approaches the point $(x_0, 0) \in \mathbb{R} \times \{0\}$ in the plane \mathbb{R}^2 , while maintaining $t > 0$. In the second limit, we take $x = x_0$ in the integrand and look at the limit $t \rightarrow 0$, still maintaining $t > 0$. Note that (48) is a 2-dimensional limit, but (49) is just a 1-dimensional limit.

Proof. Let

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy, \quad (x, t) \in \mathbb{R} \times (0, \infty). \quad (50)$$

For any $\varepsilon > 0$, we choose $\delta > 0$ such that $|\phi(y) - \phi(x_0)| < \varepsilon$ if $|y - x_0| < 2\delta$. Let $M = \sup_{\mathbb{R}} |\phi|$. If $|x - x_0| < \delta$, then

$$\begin{aligned}
|u(x, t) - \phi(x_0)| &= \left| \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} (\phi(y) - \phi(x_0)) dy \right| \\
&\leq \frac{1}{\sqrt{4\pi t}} \left(\int_{|y-x| < \delta} e^{-\frac{(x-y)^2}{4t}} |\phi(y) - \phi(x_0)| dy + \int_{|y-x| \geq \delta} e^{-\frac{(x-y)^2}{4t}} |\phi(y) - \phi(x_0)| dy \right) \\
&\leq \frac{1}{\sqrt{4\pi t}} \left(\int_{|y-x_0| < 2\delta} e^{-\frac{(x-y)^2}{4t}} |\phi(y) - \phi(x_0)| dy + 2M \int_{|y-x| \geq \delta} e^{-\frac{|x-y|^2}{4t}} dy \right) \\
&\leq \varepsilon \left(\int_{|y-x_0| < 2\delta} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy \right) + 2M \left(\int_{|y-x| \geq \delta} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy \right) \tag{51}
\end{aligned}$$

Therefore, by (42) and (46), if $t > 0$ is small enough and $|x - x_0| < \delta$, (51) will imply

$$|u(x, t) - \phi(x_0)| \leq \varepsilon + 2M\varepsilon,$$

Hence we have

$$\lim_{(x,t) \rightarrow (x_0, 0^+)} u(x, t) = \phi(x_0)$$

and (48) is proved. (49) is a consequence of (48). \square

Lemma 3.26 *Let $\phi(y)$ be a **continuous bounded** function defined on $(-\infty, \infty)$. Then we have*

$$\begin{aligned}
&\left(\frac{\partial^{m+n}}{\partial t^m \partial x^n} \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy \right) (x_0, t_0) \\
&= \int_{-\infty}^{\infty} \left[\left(\frac{\partial^{m+n}}{\partial t^m \partial x^n} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) \right) (x_0, t_0, y) \right] dy \tag{52}
\end{aligned}$$

for all $(x_0, t_0) \in (-\infty, \infty) \times (0, \infty)$ and all $m, n \in \mathbb{N} \cup \{0\}$. In particular, the function

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy, \quad (x, t) \in (-\infty, \infty) \times (0, \infty), \tag{53}$$

satisfies

$$\begin{cases} (1). u(x, t) \in C^\infty((-\infty, \infty) \times (0, \infty)) \\ (2). u_t(x, t) = u_{xx}(x, t), \quad \forall (x, t) \in (-\infty, \infty) \times (0, \infty). \end{cases} \tag{54}$$

Remark 3.27 (Important.) *To understand the proof of Lemma 3.26, you need to know when a differentiation (say $\frac{\partial}{\partial x}$) and an improper integral (say of the form $\int_{-\infty}^{\infty} g(x, y) dy$ or $\int_0^{\infty} g(x, y) dy$ for some differentiable function $g(x, y)$) can **commute**. For your convenience, here I provide two results in the following:*

1. Let $f(x, y) \in C^0(I \times [0, \infty))$, where $I \subseteq \mathbb{R}$ is an **arbitrary connected interval** and assume that the improper integral $\int_0^{\infty} f(x, y) dy$ **converges uniformly to a function $F(x)$ on I** . Then $F(x)$ is continuous on I . This means that we have the identity

$$\lim_{x \rightarrow x_0} \int_0^{\infty} f(x, y) dy = \int_0^{\infty} f(x_0, y) dy, \quad \forall x_0 \in I. \tag{55}$$

The same conclusion holds if we replace $\int_0^{\infty} f(x, y) dy$ by $\int_{-\infty}^{\infty} f(x, y) dy$.

2. Let $f(x, y) \in C^0(I \times [0, \infty))$, where $I \subseteq \mathbb{R}$ is an **arbitrary connected interval** and assume that the improper integral $\int_0^\infty f(x, y) dy$ converges to a function $F(x)$ on I (**does not have to be uniform**) and $\frac{\partial f}{\partial x} \in C^0(I \times [0, \infty))$ and $\int_0^\infty \frac{\partial f}{\partial x}(x, y) dy$ **converges uniformly** on I , Then $F(x)$ is differentiable with respect to $x \in I$ and

$$F'(x) = \int_0^\infty \frac{\partial f}{\partial x}(x, y) dy, \quad \forall x \in I. \quad (56)$$

In particular, $F(x)$ is also continuous on I . Moreover, if I is a **finite interval**, then $\int_0^\infty f(x, y) dy$ also **converges uniformly** on I . The same conclusion holds if we replace $\int_0^\infty f(x, y) dy$ by $\int_{-\infty}^\infty f(x, y) dy$ and $\int_0^\infty \frac{\partial f}{\partial x}(x, y) dy$ by $\int_{-\infty}^\infty \frac{\partial f}{\partial x}(x, y) dy$.

Note: Compare with Rudin's Advanced Calculus book (Principle of Mathematical Analysis, 3rd edition) Theorem 7.17 in p. 152. In terms of series of functions, Rudin's Theorem 7.17 can be stated as: Let $\{f_n\}$ be a sequence of differentiable functions on $[a, b]$ such that the series $\sum_{n=1}^\infty f_n(x)$ converges for some $x_0 \in [a, b]$ and assume that the series $\sum_{n=1}^\infty f'_n(x)$ converges uniformly on $[a, b]$ to a function $h(x)$, then the series $\sum_{n=1}^\infty f_n(x)$ converges uniformly on $[a, b]$ to a function $f(x)$, which is differentiable, and we have

$$f'(x) = h(x), \quad \forall x \in [a, b].$$

Proof. For any fixed $m, n \in \mathbb{N} \cup \{0\}$ and fixed $(x_0, t_0) \in (-\infty, \infty) \times (0, \infty)$ the function

$$\left(\frac{\partial^{m+n}}{\partial t^m \partial x^n} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) \right) (x_0, t_0, y), \quad y \in (-\infty, \infty)$$

decays exponentially in the variable y as $|y| \rightarrow \infty$. In fact, it also decays exponentially in the variable y as $|y| \rightarrow \infty$ for all (x, t) in some neighborhood R of (x_0, t_0) . For example, one can take R as

$$R = \left\{ (x, t) : x_0 - 1 < x < x_0 + 1, \frac{t_0}{2} < t < \frac{3t_0}{2} \right\}, \quad t_0 > 0. \quad (57)$$

By this decay property, one can check that the integral

$$\int_{-\infty}^\infty \left[\left(\frac{\partial^{m+n}}{\partial t^m \partial x^n} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) \right) (x, t, y) \right] dy, \quad (x, t) \text{ is near } (x_0, t_0)$$

converges uniformly for all (x, t) in R . By standard theory in advanced calculus, the function (as a function of $(x, t) \in R$)

$$\int_{-\infty}^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy$$

is **differentiable** with respect to t up to m times and differentiable with respect to x up to n times, i.e. one can apply $\frac{\partial^{m+n} u}{\partial t^m \partial x^n}$ onto it and obtain the identity

$$\begin{aligned} & \left(\frac{\partial^{m+n} u}{\partial t^m \partial x^n} \int_{-\infty}^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy \right) (x, t) \\ &= \int_{-\infty}^\infty \left[\left(\frac{\partial^{m+n} u}{\partial t^m \partial x^n} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) \right) (x, t, y) \right] dy, \quad \forall (x, t) \in R. \end{aligned} \quad (58)$$

As the point $(x_0, t_0) \in (-\infty, \infty) \times (0, \infty)$ is arbitrary and the numbers $m, n \in \mathbb{N} \cup \{0\}$ are also arbitrary, the identity (52) is proved for all $(x_0, t_0) \in (-\infty, \infty) \times (0, \infty)$. Moreover, the function

$$\int_{-\infty}^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy$$

is a C^∞ function of $(x, t) \in (-\infty, \infty) \times (0, \infty)$, which implies $u(x, t) \in C^\infty((-\infty, \infty) \times (0, \infty))$. Finally, we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right) u(x, t) &= \left(\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right) \left(\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy \right) \\ &= \int_{-\infty}^{\infty} \left(\left(\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right) \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \right) \phi(y) dy = \int_{-\infty}^{\infty} 0 \cdot \phi(y) dy = 0, \end{aligned}$$

which means that $u(x, t)$ satisfies the heat equation on $(-\infty, \infty) \times (0, \infty)$. \square

3.4 Heat equation on the whole line with initial condition.

Motivated by the heat polynomials, to get unique solution (we hope so) of a heat equation $u_t = u_{xx}$, we focus on the following **initial value problem**:

$$\begin{cases} u_t(x, t) = u_{xx}(x, t), & (x, t) \in \mathbb{R} \times (0, \infty) \\ u(x, 0) = f(x), & x \in \mathbb{R}. \end{cases} \quad (59)$$

Here $f(x)$ is a given **continuous function** on \mathbb{R} and we want (hope) to find a "**unique**" solution $u(x, t)$ lying in the space $C^2(\mathbb{R} \times (0, \infty)) \cap C^0(\mathbb{R} \times [0, \infty))$.

Remark 3.28 *In physical reality, most phenomena described by heat equation (and wave equation) has **initial-boundary** conditions (where the space domain for x is **bounded**). However, for $x \in (-\infty, \infty)$, the initial value problem (59) has a nice solution formula (this is similar to the wave equation $u_{tt}(x, t) = u_{xx}(x, t)$ with initial conditions $u(x, 0)$ and $u_t(x, 0)$) and it is easier to manipulate. Therefore, **for mathematical reason (not for physical reason)**, instead of looking at initial-boundary value problem for heat equation, we look at (59) first.*

Note that, unlike the wave equation, here we do not need the condition $u_t(x, 0) = g(x)$ for the heat equation. This is due to **physical phenomenon** (heat equation is not a mechanical equation coming from Newton's law) and also due to the fact that if $u(x, t)$ is C^2 up to $t = 0$ (with $f \in C^2(\mathbb{R})$), then we also have

$$u_t(x, 0) = u_{xx}(x, 0) = f''(x), \quad x \in \mathbb{R},$$

i.e. the condition $u_t(x, 0) = g(x)$ is automatically a consequence of the condition $u(x, 0) = f(x)$. On the other hand, for the case of wave equation, we can not determine $u_t(x, 0)$ from the condition $u(x, 0)$ (but we can determine $u_{tt}(x, 0)$ from the condition $u(x, 0)$ due to the identity $u_{tt}(x, 0) = u_{xx}(x, 0)$).

Unfortunately, the initial value problem (59) has infinitely many solutions (this is unlike the wave equation, which has a unique solution once we know $u(x, 0)$ and $u_t(x, 0)$) unless we impose condition on the behavior of solution $u(x, t)$ for large $|x|$. This is because the data is prescribed on the line $t = 0$, which is a **characteristic line** of the heat equation $u_{xx}(x, t) - u_t(x, t) = 0$.

In spite of this defect, when $f(x)$ is given and $x \in (-\infty, \infty)$, there is some "**special solution**" of (59), which is given by a **representation formula** (solution formula), which has **good properties** and **is close to the physical reality**.

Remark 3.29 *Recall that for a second order linear **parabolic** equation with constant coefficients for $u(x, y)$ (here we view y as time), given by*

$$au_{xx} + 2bu_{xy} + cu_{yy} + (\text{lower order terms}) = 0, \quad ac = b^2, \quad (60)$$

where a, b, c are constants with $ac = b^2$, the leading terms $au_{xx} + 2bu_{xy} + cu_{yy}$ can be factored as

$$au_{xx} + 2bu_{xy} + cu_{yy} = \left(A \frac{\partial}{\partial x} + B \frac{\partial}{\partial y} \right) \left[\left(A \frac{\partial}{\partial x} + B \frac{\partial}{\partial y} \right) u \right] = 0 \quad (61)$$

for some constants A, B, C . The **1-parameter family of lines**

$$Bx - Ay = \lambda, \quad \lambda \in (-\infty, \infty) \quad (62)$$

are called the **characteristic lines** of the parabolic equation (60). By this, for the standard heat equation $u_{xx} + (-u_t) = 0$, the **1-parameter family** of characteristic lines are given by (we now have $B = 0, A = 1, y = t$ in (62))

$$-t = \lambda, \quad \lambda \in (-\infty, \infty). \quad (63)$$

From it, we know that the line $t = 0$ (i.e. x -axis) is a **characteristic line** of the heat equation. This may explain the **nonuniqueness** of the initial value problem (59).

We now consider the following initial value problem for heat equation defined on the whole line:

$$\begin{cases} u_t = u_{xx}, & x \in (-\infty, \infty), \quad t \in (0, \infty) \\ u(x, 0) = \phi(x), & x \in (-\infty, \infty). \end{cases} \quad (64)$$

Here $\phi(x)$ is a given **continuous** function on $(-\infty, \infty)$ and we want to find a solution lying in the function space:

$$u(x, t) \in C^2((-\infty, \infty) \times (0, \infty)) \cap C^0((-\infty, \infty) \times [0, \infty)), \quad (65)$$

where satisfies (64).

As a consequence of Lemma 3.24 and Lemma 3.26, we can obtain the following **solution formula** for the initial value problem (64):

Theorem 3.30 Assume $\phi(x)$ is a **continuous bounded** function defined on $(-\infty, \infty)$. Then the function

$$u(x, t) = \begin{cases} \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy, & x \in (-\infty, \infty), \quad t \in (0, \infty) \\ \phi(x), & t = 0 \end{cases} \quad (66)$$

belongs to the space $C^\infty(\mathbb{R} \times (0, \infty)) \cap C^0(\mathbb{R} \times [0, \infty))$ (i.e. **continuous up to $t = 0$**) and satisfies the initial value problem

$$\begin{cases} u_t(x, t) = u_{xx}(x, t), & x \in (-\infty, \infty), \quad t \in (0, \infty) \\ u(x, 0) = \phi(x), & x \in (-\infty, \infty). \end{cases} \quad (67)$$

Proof. This is a direct consequence of Lemma 3.24 and Lemma 3.26. □

Remark 3.31 (Important.) As long as $t > 0$, $u(x, t)$ becomes a **smooth** function even if the initial data $\phi(x)$ is only a continuous function. We call this a **smoothing effect** of the heat equation. This is unlike the wave equation, which has no smoothing effect.

Corollary 3.32 (The maximum principle.) The solution $u(x, t)$ given by (66), where $\phi(x)$ is a **continuous bounded** function defined on $(-\infty, \infty)$, satisfies the maximum principle:

$$\inf_{\mathbb{R}} \phi \leq u(x, t) \leq \sup_{\mathbb{R}} \phi \quad \text{for all } x \in (-\infty, \infty), \quad t \in (0, \infty). \quad (68)$$

Proof. We have

$$\begin{aligned} u(x, t) &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \underbrace{\phi(y)}_{\sup_{\mathbb{R}} \phi} dy \leq \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \underbrace{\sup_{\mathbb{R}} \phi}_{\sup_{\mathbb{R}} \phi} dy \\ &= \sup_{\mathbb{R}} \phi \cdot \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy = \sup_{\mathbb{R}} \phi \end{aligned}$$

and similarly $u(x, t) \geq \inf_{\mathbb{R}} \phi$. □

Corollary 3.33 (*Infinite speed of propagation of the heat equation.*) Let $\phi(x)$ be a **continuous bounded** function defined on $(-\infty, \infty)$. Assume $\phi(x) \geq 0$ everywhere, has **compact support**, and $\phi \not\equiv 0$. Then the solution $u(x, t)$ given by (66) satisfies

$$u(x, t) > 0, \quad \forall x \in (-\infty, \infty), \quad t \in (0, \infty), \quad (69)$$

i.e., as long as time is positive, $u(x, t)$ is positive everywhere no matter how large $|x|$ is (that is why we say the equation has **infinite speed of propagation**).

Remark 3.34 This is different from the wave equation. The function $u(x, t) = \phi(x - t)$ satisfies the wave equation $u_{tt} = u_{xx}$ with $u(x, 0) = \phi(x)$. However, for $t > 0$, $u(x, t) = 0$ if $|x| > 0$ is large enough.

Proof. Since ϕ is not a zero function, we have $\phi(x_0) > 0$ for some $x_0 \in (-\infty, \infty)$. By continuity, $\phi > 0$ on $(x_0 - \varepsilon, x_0 + \varepsilon)$ for some $\varepsilon > 0$. Now at any $(x, t) \in (-\infty, \infty) \times (0, \infty)$, we have

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy \geq \int_{x_0 - \varepsilon}^{x_0 + \varepsilon} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy > 0.$$

The proof is done. □

To go on, we need the following special case of Fubini Theorem from **advanced calculus** textbook:

Lemma 3.35 (*Tonelli's theorem.*) Let $\phi(x, y)$ be a **continuous "nonnegative"** function defined on $\mathbb{R}^2 = (-\infty, \infty) \times (-\infty, \infty)$. Then the finiteness of any one of the following three integrals:

$$\int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \phi(x, y) dx \right) dy, \quad \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \phi(x, y) dy \right) dx, \quad \iint_{\mathbb{R}^2} \phi(x, y) dx dy$$

implies that of the other two. Moreover, their values are all equal.

Remark 3.36 The condition $\phi(x, y) \geq 0$ on \mathbb{R}^2 is essential.

Proof. We omit it. □

Lemma 3.37 (*Conservation of total energy.*) Let $\phi(x)$ be a **continuous bounded** function defined on $(-\infty, \infty)$ ($\phi(x)$ may not be nonnegative). Assume $\int_{-\infty}^{\infty} |\phi(x)| dx$ converges. Then the solution $u(x, t)$ given by (66) satisfies

$$\int_{-\infty}^{\infty} u(x, t) dx = \int_{-\infty}^{\infty} \phi(x) dx, \quad \forall t \in (0, \infty). \quad (70)$$

This means that **the total energy (heat) is conserved**.

Proof. For each $x \in (-\infty, \infty)$, let $\phi^+(x) = \max\{\phi(x), 0\} \geq 0$ and $\phi^-(x) = -\min\{\phi(x), 0\} \geq 0$. Then we have

$$\phi(x) = \phi^+(x) - \phi^-(x), \quad |\phi(x)| = \phi^+(x) + \phi^-(x), \quad \forall x \in (-\infty, \infty).$$

The convergence of $\int_{-\infty}^{\infty} |\phi(x)| dx$ implies that of $\int_{-\infty}^{\infty} \phi^+(x) dx$ and $\int_{-\infty}^{\infty} \phi^-(x) dx$. Also, since $\phi(x)$ is a bounded function, for each fixed $(x, t) \in (-\infty, \infty) \times (0, \infty)$, the three improper integrals

$$\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy, \quad \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi^+(y) dy, \quad \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi^-(y) dy$$

all converge. Now we have

$$\begin{aligned} \int_{-\infty}^{\infty} u(x, t) dx &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy \right] dx \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi^+(y) dy - \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi^-(y) dy \right] dx \end{aligned} \quad (71)$$

and by Lemma 3.35, we have

$$\begin{aligned} &\int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi^+(y) dy \right] dx \\ &= \int_{-\infty}^{\infty} \left[\phi^+(y) \underbrace{\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dx}_{=1} \right] dy = \int_{-\infty}^{\infty} \phi^+(y) dy < \infty \end{aligned}$$

and similarly

$$\int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi^-(y) dy \right] dx = \int_{-\infty}^{\infty} \phi^-(y) dy < \infty.$$

Therefore, the two iterated integrals in (71) converge and we conclude

$$\int_{-\infty}^{\infty} u(x, t) dx = \int_{-\infty}^{\infty} \phi^+(y) dy - \int_{-\infty}^{\infty} \phi^-(y) dy = \int_{-\infty}^{\infty} \phi(x) dx, \quad \forall t \in (0, \infty).$$

The proof is done. \square

3.5 The maximum principle.

Assume that $u(x, t)$ is a solution of the heat equation $u_t = u_{xx}$ on $(-\infty, \infty) \times (-\infty, \infty)$. The maximum principle of the diffusion equation says that (roughly speaking), for fixed time t_0 , we have $u_t(x_0, t_0) \leq 0$ if $u(x_0, t_0)$ has a *local maximum* at $x = x_0$ (so the value of $u(x_0, t_0)$ will "decrease" at the moment $t = t_0$); and $u_t(x_0, t) \geq 0$ if $u(x_0, t)$ has a *local minimum* at $x = x_0$ (so the value of $u(x_0, t_0)$ will "increase" at the moment $t = t_0$) (draw a picture for this). This is called **the maximum principle** of the heat equation. It matches with the physical phenomenon that heat goes from hot points to cold points and vice versa.

The maximum principle on the unbounded domain $x \in (-\infty, \infty)$ is more difficult to describe. We will discuss the maximum principle on **bounded domains** only.

3.6 The maximum principle on bounded domains.

Let $U_T \subset \mathbb{R}^2$ be the set given by

$$U_T = \{(x, t) \in \mathbb{R}^2 : 0 < x < \ell, 0 < t \leq T\}, \quad \ell, T > 0 \quad (72)$$

and assume that $u = u(x, t) \in C^2(U_T) \cap C^0(\bar{U}_T)$ satisfies the heat equation $u_t = u_{xx}$ on U_T (note that the segment (x, T) , $0 < x < \ell$, is included). Note that since u is continuous on the compact set \bar{U}_T , it has global maximum and minimum on \bar{U}_T .

Remark 3.38 Explain the meaning of $u \in C^2(U_T)$.

The maximum principle says the following:

Lemma 3.39 (*Weak maximum principle for heat equation.*) Assume $u \in C^2(U_T) \cap C^0(\bar{U}_T)$ satisfies the heat equation $u_t = u_{xx}$ on U_T . Then we have

$$\max_{\bar{U}_T} u = \max_{\Gamma_T} u, \quad (73)$$

where $\Gamma_T := \bar{U}_T - U_T$, which is called the **parabolic boundary** of U_T .

Remark 3.40 The above result is still true if we have $u_t \leq u_{xx}$ on U_T .

Proof. Assume $v \in C^2(U_T) \cap C^0(\bar{U}_T)$ is a function such that

$$v_{xx}(x, t) - v_t(x, t) > 0 \quad \text{in } U_T. \quad (74)$$

Then since $v \in C^0(\bar{U}_T)$, there is a point $(x_0, t_0) \in \bar{U}_T$ such that $v(x_0, t_0) = \max_{\bar{U}_T} v$. If $(x_0, t_0) \in U_T$ with $t < T$, then from calculus we know that

$$v_x(x_0, t_0) = 0, \quad v_{xx}(x_0, t_0) \leq 0, \quad v_t(x_0, t_0) = 0. \quad (75)$$

This contradicts $v_{xx} - v_t > 0$ in U_T .

If $(x, t) \in U_T$ with $t = T$, then we replace $v_t(x_0, t_0) = 0$ by $v_t(x_0, t_0) \geq 0$ in (75) and get the same contradiction. Thus the point (x_0, t_0) **must lie on the parabolic boundary** of U_T and cannot lie on U_T (for $v(x, t)$ satisfying the differential inequality (74)). In such a case we have

$$\max_{\bar{U}_T} v = \max_{\Gamma_T} v \quad (\text{call this value } M), \quad \text{where } v_{xx}(x, t) - v_t(x, t) > 0 \text{ in } U_T, \quad (76)$$

and moreover, $v(x, t)$ cannot attain the value M on U_T .

Now let $v(x, t) = u(x, t) + \varepsilon x^2$ ($\varepsilon > 0$ is a small constant), where $u \in C^2(U_T) \cap C^0(\bar{U}_T)$ satisfies the heat equation on U_T . We now have

$$v_{xx}(x, t) - v_t(x, t) = u_{xx}(x, t) + 2\varepsilon - u_t(x, t) = 2\varepsilon > 0 \quad \text{in } U_T.$$

By the above discussion, we know that

$$\max_{\bar{U}_T} v = \max_{\Gamma_T} v = \max_{\Gamma_T} (u(x, t) + \varepsilon x^2) \leq \left(\max_{\Gamma_T} u(x, t) \right) + \varepsilon \ell^2$$

and by $u(x, t) = v(x, t) - \varepsilon x^2 \leq v(x, t)$, we get

$$\max_{\bar{U}_T} u \leq \max_{\bar{U}_T} v \leq \left(\max_{\Gamma_T} u(x, t) \right) + \varepsilon \ell^2. \quad (77)$$

As $\varepsilon > 0$ is arbitrary, letting $\varepsilon \rightarrow 0^+$ (note that here ℓ is finite), we obtain $\max_{\bar{U}_T} u \leq \max_{\Gamma_T} u(x, t)$. On the other hand, we also have $\max_{\bar{U}_T} u \geq \max_{\Gamma_T} u(x, t)$. Hence $\max_{\bar{U}_T} u = \max_{\Gamma_T} u(x, t)$. \square

Exercise 3.41 Instead of using $v(x, t) = u(x, t) + \varepsilon x^2$, now use the function $v(x, t) = u(x, t) - \varepsilon t$, $\varepsilon > 0$, and repeat the same argument of proof. Can you obtain the same result?

Remark 3.42 (*Be careful.*) In the above proof, we do not exclude the possibility that the maximum of $u(x, t)$ (note that $u_t = u_{xx}$) can also be attained at some point in U_T . For example, when the solution $u(x, t)$ is a **constant**, then this can happen. However, this is the only case that can happen (this is the **strong maximum principle**).

We also have the following **minimum principle**:

Corollary 3.43 (*Weak minimum principle for heat equation.*) Assume $u \in C^2(U_T) \cap C^0(\bar{U}_T)$ satisfies the heat equation $u_t = u_{xx}$ on U_T . Then we have

$$\min_{\bar{U}_T} u = \min_{\Gamma_T} u. \quad (78)$$

Remark 3.44 The above result is still true if we have $u_t \geq u_{xx}$ on U_T .

Remark 3.45 Again, here we do not exclude the possibility that the minimum can be attained at some point in U_T .

Proof. The proof for the minimum case is similar by looking at $-u$ (it also satisfies the heat equation) and the identity $\max_{\bar{U}_T}(-u) = \max_{\Gamma_T}(-u)$ becomes $-\min_{\bar{U}_T} u = -\min_{\Gamma_T} u$. \square

Corollary 3.46 Assume $u \in C^2(U_T) \cap C^0(\bar{U}_T)$ satisfies the heat equation $u_t = u_{xx}$ on U_T and $u \equiv 0$ on the parabolic boundary Γ_T , then $u \equiv 0$ on \bar{U}_T .

Proof. This is a consequence of the maximum-minimum principle. \square

Example 3.47 (*Give this as an homework problem ...*) Let $u(x, t)$ be one of the following functions:

$$t + \frac{x^2}{2}, \quad e^{t+x}, \quad e^{t-x}, \quad e^{-t} \cos x, \quad e^{-t} \sin x, \quad e^t \cosh x, \quad e^t \sinh x, \quad (x, t) \in \mathbb{R}^2.$$

They all satisfy the heat equation $u_t = u_{xx}$ (note that $e^t \cosh x$ and $e^t \sinh x$ are linear combinations of e^{t+x} and e^{t-x}). Let $U_T = (0, 1) \times (0, T]$. We have

$$\left\{ \begin{array}{l} (1). \max_{\bar{U}_T} \left(t + \frac{x^2}{2} \right) = T + \frac{1}{2}, \quad \text{attained at } (1, T) \in \Gamma_T := \bar{U}_T - U_T \\ (2). \max_{\bar{U}_T} (e^{t+x}) = e^{T+1}, \quad \text{attained at } (1, T) \in \Gamma_T \\ (3). \max_{\bar{U}_T} (e^{t-x}) = e^{T-0}, \quad \text{attained at } (0, T) \in \Gamma_T \\ (4). \max_{\bar{U}_T} (e^{-t} \cos x) = e^{-0} \cos 0 = 1, \quad \text{attained at } (0, 0) \in \Gamma_T \\ (5). \max_{\bar{U}_T} (e^{-t} \sin x) = e^{-0} \sin 1 = \sin 1, \quad \text{attained at } (1, 0) \in \Gamma_T \\ (6). \max_{\bar{U}_T} (e^t \cosh x) = e^T \cosh 1, \quad \text{attained at } (1, T) \in \Gamma_T \\ (7). \max_{\bar{U}_T} (e^t \sinh x) = e^T \sinh 1, \quad \text{attained at } (1, T) \in \Gamma_T. \end{array} \right.$$

From the above, we see that each solution attains its maximum point on the parabolic boundary Γ_T . Also note that the maximum can be attained at **any** corner point (there are four of them) of Γ_T .

One can also use **Energy Method** (integral method) to prove the following (without using the maximum principle):

Lemma 3.48 Assume $u \in C^2(\bar{U}_T)$ satisfies the heat equation $u_t = u_{xx}$ on U_T and $u \equiv 0$ on the parabolic boundary Γ_T , then $u \equiv 0$ on \bar{U}_T .

Proof. Let $E(t)$, $0 \leq t \leq T$, be the quantity

$$E(t) = \frac{1}{2} \int_0^\ell u^2(x, t) dx \geq 0, \quad 0 \leq t \leq T.$$

Then $E(t)$ is a differentiable function on $[0, T]$, $E(0) = 0$, and we have

$$\begin{aligned} \frac{dE}{dt}(t) &= \int_0^\ell u(x, t) u_t(x, t) dx = \int_0^\ell u(x, t) u_{xx}(x, t) dx \\ &= \int_0^\ell \left[\left(\frac{d}{dx} [u(x, t) u_x(x, t)] \right) - (u_x(x, t))^2 \right] dx \\ &= [u(x, t) u_x(x, t)] \Big|_{x=0}^{x=\ell} - \int_0^\ell (u_x(x, t))^2 dx = - \int_0^\ell (u_x(x, t))^2 dx \leq 0. \end{aligned} \quad (79)$$

Hence we have

$$0 \leq E(t) \leq E(0) = 0, \quad \forall t \in [0, T]. \quad (80)$$

Thus $E(t) = 0$ for all time $t \in [0, T]$ and so $u \equiv 0$ on \bar{U}_T . \square

3.7 Discontinuous bounded initial data.

What happens if the initial condition $\phi(x)$ is a bounded function defined on $(-\infty, \infty)$ but discontinuous somewhere? (here we assume that $\phi(x)$ is discontinuous only at a **finite number** of points and at each discontinuous point x_0 both $\lim_{x \rightarrow x_0^+} \phi(x)$ and $\lim_{x \rightarrow x_0^-} \phi(x)$ exist).

We have the following interesting result:

Lemma 3.49 *Let $\phi(x)$ be a **bounded** function defined on $(-\infty, \infty)$ and at $x = x_0$ it is **discontinuous** and satisfies*

$$\lim_{x \rightarrow x_0^+} \phi(x) = A, \quad \lim_{x \rightarrow x_0^-} \phi(x) = B, \quad \text{where } A \neq B \quad (81)$$

Then the function

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy, \quad x \in (-\infty, \infty), \quad t \in (0, \infty)$$

lies in the space $u \in C^\infty((-\infty, \infty) \times (0, \infty))$ and satisfies the heat equation

$$u_t(x, t) = u_{xx}(x, t), \quad \forall (x, t) \in (-\infty, \infty) \times (0, \infty) \quad (82)$$

with

$$\lim_{t \rightarrow 0^+} u(x_0, t) = \lim_{t \rightarrow 0^+} \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x_0-y)^2}{4t}} \phi(y) dy = \frac{A+B}{2}. \quad (83)$$

Remark 3.50 (*Be careful.*) In general, the limit

$$\lim_{(x,t) \rightarrow (x_0, 0^+)} u(x, t) \quad (\text{note that this is not the same as } \lim_{t \rightarrow 0^+} u(x_0, t)) \quad (84)$$

does not exist (see Example 3.51 below). On the other hand, if $\phi(x)$ is **continuous** at $x = x_1$, then we have

$$\lim_{(x,t) \rightarrow (x_1, 0^+)} u(x, t) = \lim_{t \rightarrow 0^+} u(x_1, t) = \phi(x_1). \quad (85)$$

Proof. It suffices to verify (83). Let $M = \sup_{\mathbb{R}} |\phi|$ and let

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy, \quad (x, t) \in (-\infty, \infty) \times (0, \infty). \quad (86)$$

Then (let $y = x_0 + \sqrt{4ts}$)

$$u(x_0, t) = \frac{1}{\sqrt{\pi}} \int_0^{\infty} e^{-s^2} \phi(x_0 + \sqrt{4ts}) ds + \frac{1}{\sqrt{\pi}} \int_{-\infty}^0 e^{-s^2} \phi(x_0 + \sqrt{4ts}) ds. \quad (87)$$

For any $\varepsilon > 0$, there exists $\delta > 0$ such that if $x \in (x_0, x_0 + \delta)$, then $|\phi(x) - A| < \varepsilon$. Hence the first integral in (87) satisfies

$$\begin{aligned} & \left| \frac{1}{\sqrt{\pi}} \int_0^{\infty} e^{-s^2} \phi(x_0 + \sqrt{4ts}) ds - \frac{A}{2} \right| = \left| \frac{1}{\sqrt{\pi}} \int_0^{\infty} e^{-s^2} [\phi(x_0 + \sqrt{4ts}) - A] ds \right| \\ & \leq \frac{1}{\sqrt{\pi}} \int_0^{\delta/\sqrt{4t}} e^{-s^2} |\phi(x_0 + \sqrt{4ts}) - A| ds + \frac{1}{\sqrt{\pi}} \int_{\delta/\sqrt{4t}}^{\infty} e^{-s^2} |\phi(x_0 + \sqrt{4ts}) - A| ds \\ & \leq \frac{\varepsilon}{2} + 2M \cdot \frac{1}{\sqrt{\pi}} \int_{\delta/\sqrt{4t}}^{\infty} e^{-s^2} ds \end{aligned}$$

and so

$$\lim_{t \rightarrow 0^+} \frac{1}{\sqrt{\pi}} \int_0^{\infty} e^{-s^2} \phi(x_0 + \sqrt{4ts}) ds = \frac{A}{2}.$$

Similarly, we have

$$\lim_{t \rightarrow 0^+} \frac{1}{\sqrt{\pi}} \int_{-\infty}^0 e^{-s^2} \phi(x_0 + \sqrt{4ts}) ds = \frac{B}{2}.$$

The proof is done. □

Example 3.51 Let

$$\phi(x) = \begin{cases} 1, & x > 0 \\ 0, & x < 0, \end{cases}, \quad \phi(x) \text{ is not continuous at } x = 0.$$

It is a bounded function. Define the function

$$u(x, t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy = \int_0^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} dy$$

and let $y = x - \sqrt{4ts}$ to get (we will get the same result if we let $y = x + \sqrt{4ts}$)

$$\begin{aligned} u(x, t) &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{x/\sqrt{4t}} e^{-s^2} ds = \frac{1}{\sqrt{\pi}} \left(\int_{-\infty}^0 + \int_0^{x/\sqrt{4t}} \right) e^{-s^2} ds \\ &= \frac{1}{2} + \frac{1}{\sqrt{\pi}} \int_0^{x/\sqrt{4t}} e^{-s^2} ds, \quad (x, t) \in (-\infty, \infty) \times (0, \infty). \end{aligned}$$

We note that

$$\left\{ \begin{array}{l} u(x, t) \in C^{\infty}((-\infty, \infty) \times (0, \infty)) \text{ and } u_t = u_{xx} \text{ on } (-\infty, \infty) \times (0, \infty) \\ \lim_{(x,t) \rightarrow (x_0, 0^+)} u(x, t) = 1, \quad \text{if } x_0 > 0 \\ \lim_{(x,t) \rightarrow (x_0, 0^+)} u(x, t) = 0, \quad \text{if } x_0 < 0 \\ \lim_{t \rightarrow 0^+} u(0, t) = \frac{1}{2} = \frac{1+0}{2}, \\ \lim_{(x,t) \rightarrow (0, 0^+)} u(x, t) = \lim_{(x,t) \rightarrow (0, 0^+)} \left(\frac{1}{2} + \frac{1}{\sqrt{\pi}} \int_0^{x/\sqrt{4t}} e^{-s^2} ds \right) \text{ does not exist.} \end{array} \right. \quad (88)$$

The last limit in (88) does not exist is due to the fact that as $(x, t) \rightarrow (0, 0^+)$, the quantity $x/\sqrt{4t}$ can approach any possible number in $(-\infty, \infty)$. Hence the limit

$$\lim_{(x,t) \rightarrow (0,0^+)} \int_0^{x/\sqrt{4t}} e^{-s^2} ds$$

does not exist.

Example 3.52 Let the initial data $\phi(x)$ be

$$\phi(x) = \begin{cases} e^{-x}, & x \in (0, \infty) \\ 0, & x \in (-\infty, 0). \end{cases}$$

ϕ is bounded but not continuous at $x = 0$. Now we have

$$\begin{aligned} u(x, t) &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4t}} \phi(y) dy \\ &= \int_0^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2 - 2xy + y^2 + 4ty}{4t}} dy = \int_0^{\infty} \frac{1}{\sqrt{4\pi t}} e^{-\frac{[y+(2t-x)]^2}{4t}} e^{t-x} dy \quad (\text{let } y = x - 2t + \sqrt{4ts}) \\ &= \frac{1}{\sqrt{\pi}} e^{t-x} \int_{\frac{2t-x}{\sqrt{4t}}}^{\infty} e^{-s^2} ds, \quad (x, t) \in (-\infty, \infty) \times (0, \infty). \end{aligned}$$

It satisfies $u_t(x, t) = u_{xx}(x, t)$ on $(-\infty, \infty) \times (0, \infty)$ and

$$\left\{ \begin{array}{l} \lim_{(x,t) \rightarrow (x_0, 0^+)} \left(\frac{1}{\sqrt{\pi}} e^{t-x} \int_{\frac{2t-x}{\sqrt{4t}}}^{\infty} e^{-s^2} ds \right) = \frac{1}{\sqrt{\pi}} e^{-x_0} \int_{-\infty}^{\infty} e^{-s^2} ds = e^{-x_0}, \quad \text{if } x_0 > 0 \\ \lim_{(x,t) \rightarrow (x_0, 0^+)} \left(\frac{1}{\sqrt{\pi}} e^{t-x} \int_{\frac{2t-x}{\sqrt{4t}}}^{\infty} e^{-s^2} ds \right) = \frac{1}{\sqrt{\pi}} e^{-x_0} \int_{\infty}^{\infty} e^{-s^2} ds = 0, \quad \text{if } x_0 < 0 \\ \lim_{t \rightarrow 0^+} u(0, t) = \lim_{t \rightarrow 0^+} \left(\frac{1}{\sqrt{\pi}} e^t \int_{\frac{2t}{\sqrt{4t}}}^{\infty} e^{-s^2} ds \right) = \frac{1}{\sqrt{\pi}} \int_0^{\infty} e^{-s^2} ds = \frac{1}{2}. \end{array} \right.$$

3.8 Even, odd, and periodic initial data ($n = 1$).

Lemma 3.53 (*Even initial data for $n = 1$.*) Assume $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and is an even function with respect to $x = 0$, i.e.

$$f(-x) = f(x), \quad \forall x \in \mathbb{R}$$

Let

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} f(y) dy, \quad x \in \mathbb{R}, \quad t > 0. \quad (89)$$

Then for each $t > 0$ we have

$$u(-x, t) = u(x, t), \quad \forall x \in \mathbb{R}, \quad t > 0 \quad (90)$$

and moreover,

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_0^{\infty} \left[e^{-\frac{(x-y)^2}{4t}} + e^{-\frac{(x+y)^2}{4t}} \right] f(y) dy, \quad \forall x \in \mathbb{R}, \quad t > 0. \quad (91)$$

In particular, we have

$$u_x(0, t) = 0, \quad \forall t > 0. \quad (92)$$

Proof. For $x \in \mathbb{R}$ we have

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_0^\infty e^{-\frac{(x-y)^2}{4t}} f(y) dy + \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^0 e^{-\frac{(x-y)^2}{4t}} f(y) dy,$$

where

$$\frac{1}{\sqrt{4\pi t}} \int_{-\infty}^0 e^{-\frac{(x-y)^2}{4t}} f(y) dy = \frac{1}{\sqrt{4\pi t}} \int_0^\infty e^{-\frac{(x+z)^2}{4t}} f(-z) dz = \frac{1}{\sqrt{4\pi t}} \int_0^\infty e^{-\frac{(x+z)^2}{4t}} f(z) dz.$$

Hence we get (91). By it, we also have $u(-x, t) = u(x, t)$ for all $x \in \mathbb{R}$, $t > 0$. \square

Lemma 3.54 (*Odd initial data for $n = 1$.*) Assume $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and is an odd function with respect to $x = 0$, i.e.

$$f(-x) = -f(x), \quad \forall x \in \mathbb{R}; \quad f(0) = 0.$$

Let

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^\infty e^{-\frac{(x-y)^2}{4t}} f(y) dy, \quad x \in \mathbb{R}, \quad t > 0. \quad (93)$$

Then for each $t > 0$ we have

$$u(0, t) = 0, \quad \forall t > 0 \quad (94)$$

and

$$u(-x, t) = -u(x, t), \quad \forall x \in \mathbb{R}, \quad t > 0 \quad (95)$$

and moreover,

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_0^\infty \left[e^{-\frac{(x-y)^2}{4t}} - e^{-\frac{(x+y)^2}{4t}} \right] f(y) dy, \quad \forall x \in \mathbb{R}, \quad t > 0. \quad (96)$$

Proof. Since $f(x)$ is odd, we first have

$$u(0, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^\infty e^{-\frac{y^2}{4t}} f(y) dy = 0.$$

For $x \in \mathbb{R}$ we have

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_0^\infty e^{-\frac{(x-y)^2}{4t}} f(y) dy + \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^0 e^{-\frac{(x-y)^2}{4t}} f(y) dy,$$

where

$$\frac{1}{\sqrt{4\pi t}} \int_{-\infty}^0 e^{-\frac{(x-y)^2}{4t}} f(y) dy = \frac{1}{\sqrt{4\pi t}} \int_0^\infty e^{-\frac{(x+z)^2}{4t}} f(-z) dz = -\frac{1}{\sqrt{4\pi t}} \int_0^\infty e^{-\frac{(x+z)^2}{4t}} f(z) dz.$$

Thus we have (96), and (95) follows. \square

Remark 3.55 (*Decomposition into even solution and odd solution.*) For any initial data $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$, we can write it as

$$f(x) = \frac{f(x) + f(-x)}{2} \text{ (even function)} + \frac{f(x) - f(-x)}{2} \text{ (odd function)} := f_e(x) + f_o(x),$$

and then we can decompose $u(x, t)$ as

$$\begin{aligned} u(x, t) &= \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^\infty e^{-\frac{(x-y)^2}{4t}} f(y) dy \\ &= \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^\infty e^{-\frac{(x-y)^2}{4t}} f_e(y) dy + \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^\infty e^{-\frac{(x-y)^2}{4t}} f_o(y) dy \\ &:= u_e(x, t) + u_o(x, t), \quad x \in \mathbb{R}, \quad t > 0, \end{aligned}$$

where, for each $t > 0$, $u_e(x, t)$ is **even** in x and $u_o(x, t)$ is **odd** in x with

$$(u_e)_x(0, t) = 0 \quad \text{and} \quad u_o(0, t) = 0, \quad t > 0. \quad (97)$$

Note that both $u_e(x, t)$ and $u_o(x, t)$ are solutions to the heat equation.

Remark 3.56 (Interesting example.) In Lemma 3.54, let us take

$$f(x) = \begin{cases} 1, & x > 0, \\ -1, & x < 0, \end{cases} \quad f(0) \text{ can be any number,}$$

which is an odd function with respect to $x = 0$ and by (96) we have

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} f(y) dy = \frac{1}{\sqrt{4\pi t}} \int_0^{\infty} \left[e^{-\frac{(x-y)^2}{4t}} - e^{-\frac{(x+y)^2}{4t}} \right] dy, \quad \forall x \in \mathbb{R}, \quad t > 0.$$

For fixed $t \in (0, \infty)$ we have

$$\begin{aligned} \lim_{x \rightarrow \infty} u(x, t) &= \lim_{x \rightarrow \infty} \frac{1}{\sqrt{4\pi t}} \left(\int_0^{\infty} e^{-\frac{(x-y)^2}{4t}} dy - \int_0^{\infty} e^{-\frac{(x+y)^2}{4t}} dy \right) \\ &= \lim_{x \rightarrow \infty} \frac{1}{\sqrt{4\pi t}} \left(\int_0^{\infty} e^{-\frac{(y-x)^2}{4t}} dy - \int_0^{\infty} e^{-\frac{(y+x)^2}{4t}} dy \right) = \lim_{x \rightarrow \infty} \frac{1}{\sqrt{\pi}} \left(\int_{-\frac{x}{\sqrt{4t}}}^{\infty} e^{-z^2} dz - \int_{\frac{x}{\sqrt{4t}}}^{\infty} e^{-z^2} dz \right) = 1 \end{aligned}$$

and

$$\lim_{x \rightarrow -\infty} u(x, t) = \lim_{x \rightarrow -\infty} \frac{1}{\sqrt{\pi}} \left(\int_{-\frac{x}{\sqrt{4t}}}^{\infty} e^{-z^2} dz - \int_{\frac{x}{\sqrt{4t}}}^{\infty} e^{-z^2} dz \right) = -1.$$

The **two-dimensional limit**

$$\begin{aligned} \lim_{(x,t) \rightarrow (0,0^+)} u(x, t) &= \lim_{(x,t) \rightarrow (0,0^+)} \frac{1}{\sqrt{\pi}} \left(\int_{-\frac{x}{\sqrt{4t}}}^{\infty} e^{-z^2} dz - \int_{\frac{x}{\sqrt{4t}}}^{\infty} e^{-z^2} dz \right) \\ &= \lim_{(x,t) \rightarrow (0,0^+)} \frac{1}{\sqrt{\pi}} \int_{-x/\sqrt{4t}}^{x/\sqrt{4t}} e^{-z^2} dz = \text{does not exist.} \end{aligned} \quad (98)$$

Lemma 3.57 (Periodic initial data for $n = 1$.) Assume $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and satisfies the **periodic condition**:

$$f(x + L) = f(x), \quad \forall x \in \mathbb{R}, \quad (99)$$

for some $L > 0$. Let

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} f(y) dy, \quad x \in \mathbb{R}, \quad t > 0 \quad (100)$$

then for each $t > 0$, we have

$$u(x + L, t) = u(x, t), \quad \forall x \in \mathbb{R}, \quad t > 0. \quad (101)$$

Moreover, the integral

$$\int_0^L u(x, t) dx = \int_0^L f(x) dx, \quad t > 0 \quad (102)$$

is a constant independent of time.

Remark 3.58 (Important observation.) By (102), in case we have the property

$$\lim_{t \rightarrow \infty} u(x, t) = C \quad \text{uniformly in } x \in [0, L], \quad (103)$$

then the constant C is given by

$$C = \frac{1}{L} \int_0^L f(x) dx, \quad (104)$$

which is the average of the L -periodic function $f(x)$ over $[0, L]$.

Proof. We have

$$\begin{aligned} u(x+L, t) &= \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x+L-y)^2}{4t}} f(y) dy \quad (\text{let } z = y - L) \\ &= \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-z)^2}{4t}} f(z+L) dz = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-z)^2}{4t}} f(z) dz = u(x, t). \end{aligned}$$

To see (102), we compute

$$\frac{d}{dt} \int_0^L u(x, t) dx = \int_0^L u_{xx}(x, t) dx = u_x(L, t) - u_x(0, t) = 0, \quad t > 0$$

and (102) follows. \square

Assume $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ is a 2π -**periodic** function. As for each $t > 0$, $u(x, t)$ is a smooth 2π -periodic function in x , we can use **Fourier series** to express $u(x, t)$ as

$$u(x, t) = \frac{a_0(t)}{2} + \sum_{n=1}^{\infty} a_n(t) \cos nx + b_n(t) \sin nx, \quad (x, t) \in S^1 \times (0, \infty), \quad (105)$$

where

$$\begin{cases} a_n(t) = \frac{1}{\pi} \int_0^{2\pi} u(\theta, t) \cos(n\theta) d\theta, & n = 0, 1, 2, 3, \dots, \\ b_n(t) = \frac{1}{\pi} \int_0^{2\pi} u(\theta, t) \sin(n\theta) d\theta, & n = 1, 2, 3, \dots \end{cases} \quad (106)$$

and moreover, any derivative of $u(x, t)$ can commute with the summation sign. Since we have $u_t = u_{xx}$, it implies

$$\begin{aligned} &\frac{a'_0(t)}{2} + \sum_{n=1}^{\infty} a'_n(t) \cos nx + b'_n(t) \sin nx \\ &= - \sum_{n=1}^{\infty} n^2 a_n(t) \cos nx + n^2 b_n(t) \sin nx, \quad (x, t) \in S^1 \times (0, \infty) \end{aligned}$$

and we conclude ($u(x, 0) = f(x)$)

$$a_0(t) = a_0(0), \quad a_n(t) = a_n(0) e^{-n^2 t}, \quad b_n(t) = b_n(0) e^{-n^2 t}, \quad \forall t \in (0, \infty), \quad (107)$$

where

$$\begin{cases} a_n(0) = \frac{1}{\pi} \int_0^{2\pi} f(\theta) \cos(n\theta) d\theta, & n = 0, 1, 2, 3, \dots \\ b_n(0) = \frac{1}{\pi} \int_0^{2\pi} f(\theta) \sin(n\theta) d\theta, & n = 1, 2, 3, \dots \end{cases} \quad (108)$$

and by the **Riemann-Lebesgue Lemma**, we know that $a_n(0) \rightarrow 0$, $b_n(0) \rightarrow 0$ as $n \rightarrow \infty$.

We can conclude the following:

Lemma 3.59 Assume $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ is a 2π -periodic function. Then the function

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} f(y) dy, \quad x \in \mathbb{R}, \quad t > 0 \quad (109)$$

can be expressed as

$$u(x, t) = \begin{cases} \frac{1}{2\pi} \int_0^{2\pi} f(\theta) d\theta \\ + \sum_{n=1}^{\infty} e^{-n^2 t} \left[\left(\frac{1}{\pi} \int_0^{2\pi} f(\theta) \cos(n\theta) d\theta \right) \cos nx + \left(\frac{1}{\pi} \int_0^{2\pi} f(\theta) \sin(n\theta) d\theta \right) \sin nx \right] \end{cases} \quad (110)$$

for all $(x, t) \in S^1 \times (0, \infty)$.

Remark 3.60 (Important observation.) Here is a different approach. Note that for each $n = 0, 1, 2, 3, \dots$ the convolution solution of the problem

$$\begin{cases} u_t(x, t) = u_{xx}(x, t), & x \in \mathbb{R}, \quad t > 0 \\ u(x, 0) = \cos nx \quad (\sin nx), & x \in \mathbb{R} \end{cases}$$

is given by $u(x, t) = e^{-n^2 t} \cos nx$ ($e^{-n^2 t} \sin nx$). Hence we have

$$\begin{cases} \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} (\cos ny) dy = e^{-n^2 t} \cos nx \\ \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} (\sin ny) dy = e^{-n^2 t} \sin nx. \end{cases} \quad (111)$$

By (111), we have

$$\begin{aligned} & \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} f(y) dy \\ &= \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} \left(\frac{1}{2\pi} \int_0^{2\pi} f(\theta) d\theta + \sum_{n=1}^{\infty} \left(\frac{1}{\pi} \int_0^{2\pi} f(\theta) \cos(n\theta) d\theta \right) \cos ny \right. \\ & \quad \left. + \sum_{n=1}^{\infty} \left(\frac{1}{\pi} \int_0^{2\pi} f(\theta) \sin(n\theta) d\theta \right) \sin ny \right) dy \\ &= \begin{cases} \frac{1}{2\pi} \int_0^{2\pi} f(\theta) d\theta \\ + \sum_{n=1}^{\infty} e^{-n^2 t} \left[\left(\frac{1}{\pi} \int_0^{2\pi} f(\theta) \cos(n\theta) d\theta \right) \cos nx + \left(\frac{1}{\pi} \int_0^{2\pi} f(\theta) \sin(n\theta) d\theta \right) \sin nx \right], \end{cases} \end{aligned} \quad (112)$$

which is exactly (110).

Remark 3.61 (Another simple way to verify (111).) For convenience, we look at the case $n = 1$ only. We note that

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} \sin y dy = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-z^2} \sin(x + \sqrt{4t}z) dz,$$

which gives

$$u_{xx}(x, t) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-z^2} \frac{\partial^2}{\partial x^2} \sin(x + \sqrt{4t}z) dz = -u(x, t).$$

The ODE for $u(x, t)$ implies that $u(x, t)$ has the form

$$u(x, t) = A(t) \cos x + B(t) \sin x$$

for some coefficients functions $A(t)$, $B(t)$. Since $u(x, t)$ satisfies $u_t = u_{xx}$ and $u(x, 0) = \sin x$, we must have

$$A'(t) = -A(t), \quad A(0) = 0 \quad \text{and} \quad B'(t) = -B(t), \quad B(0) = 1,$$

yielding $u(x, t) = e^{-t} \sin x$. The proof for the case $u(x, 0) = \cos x$ is similar.

Our final important result in the periodic case is the following:

Lemma 3.62 *Assume $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ is a 2π -periodic function. Then the function*

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} f(y) dy, \quad x \in \mathbb{R}, \quad t > 0 \quad (113)$$

satisfies

$$\lim_{t \rightarrow \infty} u(x, t) = \frac{1}{2\pi} \int_0^{2\pi} f(\theta) d\theta \quad \text{uniformly in } x \in \mathbb{R} \text{ (or } x \in S^1) \quad (114)$$

(in fact, $u(x, t) \rightarrow \frac{1}{2\pi} \int_0^{2\pi} f(\theta) d\theta$ in $C^\infty(S^1)$ as $t \rightarrow \infty$).

Remark 3.63 *There is a higher dimensional analogue of the above lemma. Also see Lemma ??.*

3.9 Splitting the representation formula ($n = 1$).

We first note the following:

Lemma 3.64 *Assume $f \in C^0[0, \infty) \cap L^\infty[0, \infty)$. Then we have*

$$\lim_{t \rightarrow 0^+} \int_0^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{y^2}{4t}} f(y) dy = \frac{f(0)}{2}. \quad (115)$$

Proof. We first have

$$\int_0^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{y^2}{4t}} dy = \int_{-\infty}^0 \frac{1}{\sqrt{4\pi t}} e^{-\frac{y^2}{4t}} dy = \frac{1}{\sqrt{\pi}} \int_0^\infty e^{-z^2} dz = \frac{1}{2}. \quad (116)$$

One can extend $f(x)$ to the whole line by letting it equal to $f(0)$ on $(-\infty, 0)$. Then $f(x) \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and we get

$$\begin{aligned} f(0) &= \lim_{t \rightarrow 0^+} \int_{-\infty}^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{y^2}{4t}} f(y) dy = \lim_{t \rightarrow 0^+} \left(\int_{-\infty}^0 \frac{1}{\sqrt{4\pi t}} e^{-\frac{y^2}{4t}} f(0) dy + \int_0^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{y^2}{4t}} f(y) dy \right) \\ &= \frac{f(0)}{2} + \lim_{t \rightarrow 0^+} \int_0^\infty \frac{1}{\sqrt{4\pi t}} e^{-\frac{y^2}{4t}} f(y) dy. \end{aligned}$$

The result follows. □

For $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$, let

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^\infty e^{-\frac{(x-y)^2}{4t}} f(y) dy, \quad x \in \mathbb{R}, \quad t > 0. \quad (117)$$

We know that it satisfies

$$\begin{cases} (1). \quad u \in C^\infty(\mathbb{R} \times (0, \infty)) \\ (2). \quad u_t(x, t) - u_{xx}(x, t) = 0 \quad \text{in } \mathbb{R} \times (0, \infty) \\ (3). \quad \lim_{(x,t) \rightarrow (x_0, 0^+)} u(x, t) = f(x_0) \quad \text{for all } x_0 \in \mathbb{R}. \end{cases} \quad (118)$$

For a fixed number $a \in \mathbb{R}$, we let

$$u_1(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^a e^{-\frac{(x-y)^2}{4t}} f(y) dy, \quad u_2(x, t) = \frac{1}{\sqrt{4\pi t}} \int_a^\infty e^{-\frac{(x-y)^2}{4t}} f(y) dy \quad (119)$$

for $x \in \mathbb{R}$, $t > 0$. We can see that $u_1(x, t)$ and $u_2(x, t)$ also satisfy (1), (2) of (118) with

$$u(x, t) = u_1(x, t) + u_2(x, t), \quad \forall (x, t) \in \mathbb{R} \times (0, \infty)$$

and we also have the following 2-dimensional limit results (check it yourself):

$$\lim_{(x,t) \rightarrow (x_0, 0^+)} u_1(x, t) = \begin{cases} 0, & x_0 \in (a, \infty) \\ f(x_0), & x_0 \in (-\infty, a) \end{cases} \quad (120)$$

and

$$\lim_{(x,t) \rightarrow (x_0, 0^+)} u_2(x, t) = \begin{cases} f(x_0), & x_0 \in (a, \infty) \\ 0, & x_0 \in (-\infty, a). \end{cases} \quad (121)$$

Therefore, both $u_1(x, t)$ and $u_2(x, t)$ are smooth solutions to the heat equation on $\mathbb{R} \times (0, \infty)$, but with discontinuous initial data if $f(a) \neq 0$.

At $x_0 = a$, we note that

$$u_1(a, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^a e^{-\frac{(a-y)^2}{4t}} f(y) dy = \frac{1}{\sqrt{4\pi t}} \int_0^{\infty} e^{-\frac{z^2}{4t}} f(a-z) dz, \quad y = a-z$$

and by Lemma 3.64, we have (note that $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$)

$$\lim_{t \rightarrow 0^+} u_1(a, t) = \lim_{t \rightarrow 0^+} \frac{1}{\sqrt{4\pi t}} \int_0^{\infty} e^{-\frac{z^2}{4t}} f(a-z) dz = \frac{f(a)}{2} \quad (122)$$

and similarly

$$\lim_{t \rightarrow 0^+} u_2(a, t) = \frac{f(a)}{2}. \quad (123)$$

Thus we have

$$\lim_{t \rightarrow 0^+} u(a, t) = \lim_{t \rightarrow 0^+} [u_1(a, t) + u_2(a, t)] = \frac{f(a)}{2} + \frac{f(a)}{2} = f(a),$$

consistent with our known result.

However, the **two-dimensional** limit $\lim_{(x,t) \rightarrow (a, 0^+)} u_1(x, t)$ **does not** exist in general (unless $f(a) = 0$). The same for $\lim_{(x,t) \rightarrow (a, 0^+)} u_2(x, t)$. This is because

$$\begin{aligned} & \lim_{(x,t) \rightarrow (a, 0^+)} u_1(x, t) \\ &= \lim_{(x,t) \rightarrow (a, 0^+)} \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^a e^{-\frac{(x-y)^2}{4t}} f(y) dy \quad (\text{let } y = x + \sqrt{4t}z, z \in (-\infty, (a-x)/\sqrt{4t})) \\ &= \lim_{(x,t) \rightarrow (a, 0^+)} \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\frac{a-x}{\sqrt{4t}}} e^{-z^2} f(x + \sqrt{4t}z) dz, \end{aligned}$$

where the above limit **does not** exist if $f(a) \neq 0$ due to the term $(a-x)/\sqrt{4t}$, which can tend to $+\infty$, $-\infty$, or any finite number when $(x, t) \rightarrow (a, 0^+)$. However, we have

$$\lim_{(x,t) \rightarrow (a, 0^+)} [u_1(x, t) + u_2(x, t)] = \lim_{(x,t) \rightarrow (a, 0^+)} u(x, t) = f(a).$$

More precisely, we have:

Lemma 3.65 *Let $f \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$. If $f(a) \neq 0$, the **two-dimensional limit***

$$\lim_{(x,t) \rightarrow (a, 0^+)} \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\frac{a-x}{\sqrt{4t}}} e^{-z^2} f(x + \sqrt{4t}z) dz \quad (124)$$

does not exist, and if $f(a) = 0$, the above two-dimensional limit is **equal to 0** and we have

$$\lim_{(x,t) \rightarrow (a, 0^+)} \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\frac{a-x}{\sqrt{4t}}} e^{-z^2} f(x + \sqrt{4t}z) dz = f(a) (= 0). \quad (125)$$

Remark 3.66 Note that we can express $u_1(x, t)$ as

$$u_1(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^a e^{-\frac{(x-y)^2}{4t}} f(y) dy = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} F(y) dy,$$

where

$$F(y) = \begin{cases} f(y), & y \in (-\infty, a) \\ 0, & y \in (a, \infty). \end{cases} \quad (126)$$

The function $F(y)$ can be made **continuous** at $y = a$ if $f(a) = 0$ and we define $F(a) = 0$; and it has a **jump discontinuity** at $y = a$ if $f(a) \neq 0$. Therefore, if $f(a) = 0$, we have

$$\lim_{(x,t) \rightarrow (a,0^+)} u_1(x, t) = \lim_{(x,t) \rightarrow (a,0^+)} \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} F(y) dy = F(a) = 0. \quad (127)$$

Also see Example ??.

Proof. Assume $f(a) \neq 0$. As $(x, t) \rightarrow (a, 0^+)$, the quantity $(a - x)/\sqrt{4t}$ can approach any number, or $+\infty$, or $-\infty$, and the function $f(x + \sqrt{4tz})$ for bounded $|z| \leq M$, will converge to $f(a) \neq 0$. From this, one can easily see that the limit does not exist as $(x, t) \rightarrow (a, 0^+)$. On the other hand, if $f(a) = 0$, the integral

$$\int_{-M}^M e^{-z^2} f(x + \sqrt{4tz}) dz \rightarrow \int_{-M}^M e^{-z^2} f(a) dz = 0 \quad \text{as } (x, t) \rightarrow (a, 0^+).$$

Also the two integrals (for large constant $M > 0$) both have small values:

$$\int_{-\infty}^{-M} e^{-z^2} f(x + \sqrt{4tz}) dz, \quad \int_M^{\infty} e^{-z^2} f(x + \sqrt{4tz}) dz,$$

which will imply that the limit is equal to 0 as $(x, t) \rightarrow (a, 0^+)$. \square

3.10 The equivalence between the initial condition (at $t = 0$) and the boundary conditions (at $x = 0$) for $n = 1$.

Let $u(x, t)$ be a solution of the heat equation on \mathbb{R} given by

$$u(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} g(y) dy, \quad x \in \mathbb{R}, \quad t > 0 \quad (128)$$

and $v(x, t)$ be another solution of the heat equation on \mathbb{R} given by

$$v(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} e^{-\frac{(x-y)^2}{4t}} h(y) dy, \quad x \in \mathbb{R}, \quad t > 0, \quad (129)$$

where both $g, h \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$. We claim the following result:

Lemma 3.67 Let $u(x, t)$ and $v(x, t)$ be **convolution solutions** of the heat equation given by (128) and (129) respectively. If $u(x, t)$ and $v(x, t)$ satisfy the same boundary conditions, given by

$$u(0, t) = v(0, t) \quad \text{and} \quad u_x(0, t) = v_x(0, t), \quad \forall t \in (0, \infty). \quad (130)$$

Then we must have

$$g(y) = h(y), \quad \forall y \in (-\infty, \infty), \quad (131)$$

i.e. **they have the same initial condition**. As a consequence, we have $u(x, t) \equiv v(x, t)$ for all $x \in \mathbb{R}, t \in (0, \infty)$.

Proof. It suffices to show that for a function $g \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$, if we have

$$\int_{-\infty}^{\infty} e^{-\frac{y^2}{4t}} g(y) dy = 0 \quad \text{and} \quad \int_{-\infty}^{\infty} e^{-\frac{y^2}{4t}} yg(y) dy = 0, \quad \forall t \in (0, \infty), \quad (132)$$

then we must have $g(y) \equiv 0$. Since the identities in (132) are true for all t , one can differentiate the first identity with respect to t successively and get (note that ∂_t can move into the integral sign even if the function $y^{2k}g(y)$ is unbounded on \mathbb{R})

$$\int_{-\infty}^{\infty} e^{-\frac{y^2}{4t}} y^{2n} g(y) dy = 0, \quad \forall n = 0, 1, 2, 3, \dots, \quad \forall t \in (0, \infty). \quad (133)$$

Similarly, if we do the same on the second identity of (132), we get

$$\int_{-\infty}^{\infty} e^{-\frac{y^2}{4t}} y^{2n+1} g(y) dy = 0, \quad \forall n = 0, 1, 2, 3, \dots, \quad \forall t \in (0, \infty). \quad (134)$$

Thus we conclude

$$\int_{-\infty}^{\infty} \left(e^{-\frac{y^2}{4t}} g(y) \right) y^n dy = 0, \quad \forall n = 0, 1, 2, 3, \dots, \quad \forall t \in (0, \infty), \quad (135)$$

which can imply that $g(y) \equiv 0$ for all $y \in (-\infty, \infty)$ (I cannot provide the details here, but I think it must be right; it seems that one cannot use the Exercise 20 in p. 169 of Rudin's book here since **the domain of integration is not compact**). The proof is done. \square

Remark 3.68 (Important observation ...) We need *two identities* in (132) to infer $g \equiv 0$. Just one identity in (132) is **not enough** to infer $g \equiv 0$. For example, if $g(y)$ is any odd function, we always have $\int_{-\infty}^{\infty} e^{-y^2/4t} g(y) dy = 0$. Similarly, if $g(y)$ is any even function, we always have $\int_{-\infty}^{\infty} e^{-y^2/4t} yg(y) dy = 0$.

Remark 3.69 (Important observation ...) Note that the identity in (135) is valid for all $n = 0, 1, 2, 3, \dots$ and for all $t \in (0, \infty)$. In case $g(y) \in C_b^\infty(\mathbb{R})$, we can write the identity as (let $y = 2\sqrt{tz}$)

$$\int_{-\infty}^{\infty} \left(e^{-\frac{y^2}{4t}} g(y) \right) y^n dy = \left(2\sqrt{t} \right)^{n+1} \int_{-\infty}^{\infty} \left(e^{-z^2} g(2\sqrt{tz}) \right) z^n dz = 0$$

for all $n = 0, 1, 2, 3, \dots$ and for all $t \in (0, \infty)$, which is the same as

$$\int_{-\infty}^{\infty} \left(e^{-z^2} g(2\sqrt{tz}) \right) z^n dz = 0$$

for all $n = 0, 1, 2, 3, \dots$ and for all $t \in (0, \infty)$. One can differentiate the identity with respect to $t \in (0, \infty)$ (as many times as we want) and get

$$\begin{aligned} \int_{-\infty}^{\infty} e^{-z^2} g'(2\sqrt{tz}) z^{n+1} dz &= \int_{-\infty}^{\infty} e^{-z^2} g''(2\sqrt{tz}) z^{n+2} dz \\ &= \int_{-\infty}^{\infty} e^{-z^2} g'''(2\sqrt{tz}) z^{n+3} dz = \int_{-\infty}^{\infty} e^{-z^2} g^{(4)}(2\sqrt{tz}) z^{n+4} dz = \dots = 0 \end{aligned}$$

for all $n = 0, 1, 2, 3, \dots$ and for all $t \in (0, \infty)$. Thus one can produce another new identities, which can convince us that $g(y) \equiv 0$ for all $y \in (-\infty, \infty)$.

Remark 3.70 *By the second identity, we also have (assume further that g is differentiable)*

$$0 = \int_{-\infty}^{\infty} e^{-\frac{y^2}{4t}} yg(y) dy = (-2t) \int_{-\infty}^{\infty} g(y) \frac{\partial}{\partial y} \left(e^{-\frac{y^2}{4t}} \right) dy = 2t \int_{-\infty}^{\infty} e^{-\frac{y^2}{4t}} g'(y) dy, \quad \forall t \in (0, \infty),$$

which gives the extra identity

$$\int_{-\infty}^{\infty} e^{-\frac{y^2}{4t}} g'(y) dy = 0, \quad \forall t \in (0, \infty). \quad (136)$$

Remark 3.71 (*Analytic consideration; important observation ...*) *Let $u(x, t)$ be a solution of the heat equation $u_t(x, t) = u_{xx}(x, t)$ on $\mathbb{R} \times (0, \infty)$ with*

$$u(0, t) = u_x(0, t) = 0, \quad \forall t \in (0, \infty). \quad (137)$$

Then we have

$$u_t(0, t) = (u_x)_t(0, t) = 0, \quad \forall t \in (0, \infty).$$

The above implies (note that $u_t = u_{xx}$ and $(u_x)_t = (u_t)_x = u_{xxx}$)

$$u_{xx}(0, t) = u_{xxx}(0, t) = 0, \quad \forall t \in (0, \infty). \quad (138)$$

If we keep going by differentiating with respect to t , we can get

$$\frac{\partial^m u}{\partial x^m}(0, t) = 0, \quad \forall m = 0, 1, 2, 3, \dots, \quad \forall t \in (0, \infty). \quad (139)$$

Since we know that $u(x, t)$ is **analytic** in x for each fixed $t > 0$ (this property is valid for all solutions of the heat equation, not necessarily confined to convolution solutions), (139) can imply that $u(x, t) \equiv 0$ for all $x \in (-\infty, \infty)$, $t \in (0, \infty)$.

This is the end of parabolic equations.