



# A reaction–diffusion model of harmful algae and zooplankton in an ecosystem



Sze-Bi Hsu<sup>a,1</sup>, Feng-Bin Wang<sup>b,c,\*,2</sup>, Xiao-Qiang Zhao<sup>d,e,3</sup>

<sup>a</sup> Department of Mathematics, National Tsing Hua University, Hsinchu 300, Taiwan

<sup>b</sup> Department of Natural Science in the Center for General Education, Chang Gung University, Kwei-Shan, Taoyuan 333, Taiwan

<sup>c</sup> Community Medicine Research Center, Keelung Chang Gung Memorial Hospital, Keelung 204, Taiwan

<sup>d</sup> School of Mathematics, Northwest University, Xi'an, Shaanxi 710127, China

<sup>e</sup> Department of Mathematics and Statistics, Memorial University of Newfoundland, St. John's, NL A1C 5S7, Canada

## ARTICLE INFO

### Article history:

Received 22 September 2016

Available online 22 February 2017

Submitted by Y. Yamada

### Keywords:

Chemostat

Algal blooms

Inhibition

Global attractor

Weak repellers

Coexistence

## ABSTRACT

This paper is devoted to the investigation of an unstirred chemostat system modeling the interactions of two essential nutrients (i.e., nitrogen and phosphorus), harmful algae (i.e., *P. parvum* and cyanobacteria), and a small-bodied zooplankton in an ecosystem. To obtain a weakly repelling property of a compact and invariant set on the boundary, we introduce an associated elliptic eigenvalue problem. It turns out that the model system admits a coexistence steady state and is uniformly persistent provided that the trivial steady state, two semi-trivial steady states and a global attractor on the boundary are all weak repellers.

© 2017 Elsevier Inc. All rights reserved.

## 1. Introduction

Harmful algal blooms (HABs) have been a serious problem in many coastal and inland waters worldwide [4,7]. It was known that the algal species, *Prymnesium parvum* (golden algae), is responsible for fish-killing problem, and results in major economic damage [5]. In a reservoir, *P. parvum* competes for nitrogen and phosphorus with cyanobacteria, which also excrete allelopathic cyanotoxins that inhibit the growth of *P. parvum*. A small-bodied zooplankton population consume both types of algae for growth, but the dissolved toxins produced by *P. parvum* also inhibit zooplankton ingestion, growth and reproduction. In order to understand such complex interactions and reactions in an ecosystem, the authors in [6] pro-

\* Corresponding author.

E-mail addresses: fbwang@mail.cgu.edu.tw, fbwang0229@gmail.com (F.-B. Wang).

<sup>1</sup> Research supported in part by Ministry of Science and Technology, Taiwan.

<sup>2</sup> Research supported in part by Ministry of Science and Technology, Taiwan; and National Center for Theoretical Sciences (NCTS), National Taiwan University; and Chang Gung Memorial Hospital under grants BMRPD18 and NMRPD5F0541.

<sup>3</sup> Research supported in part by the NSERC of Canada.

posed a well-mixed chemostat system to explore the dynamics of nutrients, *P. parvum*, toxin(s) produced by *P. parvum*, cyanobacteria, cyanotoxin(s) produced by cyanobacteria, and zooplankton.

A natural approach to the spatial heterogeneity is to use “unstirred” chemostat, where we will remove the assumption that interactions of nutrients and species proceeds in a well-mixed, spatially uniform habitat. The unstirred chemostat can be regarded as a spatially distributed habitat in which inflow of nutrients occurs at one point and outflow at another, with diffusive transport of nutrients and organisms between these points [3,14]. For simplicity, we ignore the equations of toxins proposed in [6] and assume that inhibitory effects are directly determined by the densities of harmful algae. Accordingly, we modify the model in [6] to obtain the following unstirred chemostat model:

$$\begin{cases} \frac{\partial R}{\partial t} = d\frac{\partial^2 R}{\partial x^2} - q_{1r}f_1(R, S)u_1e^{-\alpha u_2} - q_{2r}f_2(R, S)u_2, & x \in (0, 1), t > 0, \\ \frac{\partial S}{\partial t} = d\frac{\partial^2 S}{\partial x^2} - q_{1s}f_1(R, S)u_1e^{-\alpha u_2} - q_{2s}f_2(R, S)u_2, & x \in (0, 1), t > 0, \\ \frac{\partial u_1}{\partial t} = d\frac{\partial^2 u_1}{\partial x^2} + f_1(R, S)u_1e^{-\alpha u_2} - q_1g_1(u_1)Z, & x \in (0, 1), t > 0, \\ \frac{\partial u_2}{\partial t} = d\frac{\partial^2 u_2}{\partial x^2} + f_2(R, S)u_2 - q_2g_2(u_2)Z, & x \in (0, 1), t > 0, \\ \frac{\partial Z}{\partial t} = d\frac{\partial^2 Z}{\partial x^2} + G(u_1, u_2)Z, & x \in (0, 1), t > 0, \end{cases} \tag{1.1}$$

with boundary conditions

$$\begin{cases} \frac{\partial R}{\partial x}(0, t) = -R^{(0)}, \quad \frac{\partial R}{\partial x}(1, t) + \gamma R(1, t) = 0, & t > 0, \\ \frac{\partial S}{\partial x}(0, t) = -S^{(0)}, \quad \frac{\partial S}{\partial x}(1, t) + \gamma S(1, t) = 0, & t > 0, \\ \frac{\partial u_i}{\partial x}(0, t) = \frac{\partial u_i}{\partial x}(1, t) + \gamma u_i(1, t) = 0, & t > 0, \quad i = 1, 2, \\ \frac{\partial Z}{\partial x}(0, t) = \frac{\partial Z}{\partial x}(1, t) + \gamma Z(1, t) = 0, & t > 0, \end{cases} \tag{1.2}$$

and initial conditions

$$\begin{cases} R(x, 0) = R^0(x) \geq 0, S(x, 0) = S^0(x) \geq 0, & x \in (0, 1), \\ u_i(x, 0) = u_i^0(x) \geq 0, Z(x, 0) = Z^0(x) \geq 0, & x \in (0, 1), \quad i = 1, 2. \end{cases} \tag{1.3}$$

Here  $R(x, t)$  and  $S(x, t)$  denote the complementary nutrient (nitrogen and phosphorus) concentrations at position  $x$  and time  $t$ ;  $u_1(x, t)$  and  $u_2(x, t)$  represent the densities of *P. parvum* (golden algae) and cyanobacteria, respectively;  $Z(x, t)$  represents the density of small-bodied zooplankton population.  $R^{(0)}$  and  $S^{(0)}$  are input concentration of nutrients;  $q_{ir}$  and  $q_{is}$ ,  $i = 1, 2$ , are the constant nutrient quotas;  $q_i$ ,  $i = 1, 2$ , is the constant algal quota; the constant  $\gamma$  in (1.2) represents the washout constant. We also assume that nutrients and algal species have the same diffusion coefficient  $d$ . The term  $e^{-\alpha u_2}$  describes the inhibitory effect on  $u_1(x, t)$  from  $u_2(x, t)$ . The response function are given by  $f_i(R, S) = \min\{h_{ir}(R), h_{is}(S)\}$ ,  $i = 1, 2$ . The nonlinear functions  $h_{ir}(R)$  ( $h_{is}(S)$ ) describe the nutrient uptake and growth rates of species  $i$  when only nutrient  $R$  ( $S$ ) is limiting. We assume that the functions  $h_{ir}(R)$  and  $h_{is}(S)$  satisfy

$$h_{ir}(0) = 0, h'_{ir}(R) > 0 \quad \forall R > 0, h_{ir} \in C^2, i = 1, 2.$$

An usual example is the Monod function

$$h_{ir}(R) = \frac{m_{ir}R}{K_{ir} + R}, \quad h_{is}(S) = \frac{m_{is}S}{K_{is} + S}.$$

Both types of algae are consumed by zooplankton, and consumption of the algae supports the growth of the zooplankton. Further, *P. parvum*  $u_1(x, t)$  also inhibits the growth of zooplankton. The function  $g_1(u_1)$  represents the relationship between zooplankton and *P. parvum*, which simultaneously include positive and

negative effects on the growth of the zooplankton, depending on the density of  $u_1(x, t)$ . Then  $g_1(u_1)$  takes the following form:

$$g_1(u_1) = \frac{m_{1z}u_1}{K_{1z} + u_1 + \eta u_1^2}, \quad \text{or } g_1(u_1) = \frac{m_{1z}u_1}{K_{1z} + u_1} e^{-\beta u_1}.$$

The cyanobacteria  $u_2(x, t)$  only inhibits the growth of P. parvum  $u_1(x, t)$ , and has no negative effects on zooplankton  $Z(x, t)$ . The function  $g_2(u_2)$  represents the relationship between zooplankton and cyanobacteria  $u_2(x, t)$ , and  $g_2(u_2)$  is increasing in  $u_2$ , and hence,  $g_2(u_2)$  takes the following form:

$$g_2(u_2) = \frac{m_{2z}u_2}{K_{2z} + u_2}.$$

Then  $G(u_1, u_2)$ , the growth rate of zooplankton, takes the following types

$$G(u_1, u_2) = g_1(u_1) \cdot g_2(u_2), \tag{1.4}$$

or

$$G(u_1, u_2) = \min\{g_1(u_1), g_2(u_2)\}. \tag{1.5}$$

The organization of this paper is as follows. In Section 2, we study the well-posedness of system (1.1)–(1.3). Section 3 is devoted to the study of the global dynamics of system (3.1)–(3.3) modeling the interactions of P. parvum and cyanobacteria, that is, the subsystem of (1.1)–(1.3) where we put  $Z(x, t) \equiv 0$ . Basically, we show that the semiflow generated by system (3.1)–(3.3) admits a global attractor  $A_0 \subset \text{Int}(C([0, 1], \mathbb{R}_+^4))$  when the semi-trivial steady-state solutions of (3.1)–(3.3) are both unstable, by appealing to the theory of uniform persistence and chain transitive sets. In Section 4, we investigate the coexistence of harmful algae (i.e., P. parvum and cyanobacteria) and zooplankton for system (1.1)–(1.3). The main difficulty is that the zooplankton-extinct steady-state solution of system (1.1)–(1.3) is not necessarily unique, that is, the set  $A_0 \times \{0\}$  may not be a singleton. To address this general case, we first introduce a continuous function,  $m(x)$ , involving in  $A_0$  (see (4.3)), and the principal eigenvalue of an eigenvalue problem associated with  $m(x)$  (see (4.7)) becomes a crucial index that determines whether  $M_3 := A_0 \times \{0\}$  is a uniform weak repeller in the sense of (4.8). Then we use persistence theory to establish the existence of a positive (coexistence) steady state and the uniform persistence for system (1.1)–(1.3) under the assumption that its compact invariant set  $M_3 := A_0 \times \{0\}$ , trivial and semitrivial steady states are all weak repellers. A brief discussion section completes the paper.

## 2. Well-posedness

We first study the well-posedness of the initial-boundary-value problem (1.1)–(1.3). Let  $\mathbb{X} = C([0, 1], \mathbb{R}_+^5)$  be the positive cone in the Banach space  $C([0, 1], \mathbb{R}^5)$  with the usual supremum norm. In order to simplify notations, we set  $v_1 = R$ ,  $v_2 = S$ ,  $v_3 = u_1$ ,  $v_4 = u_2$ ,  $v_5 = Z$  and  $\mathbf{v} = (v_1, v_2, v_3, v_4, v_5)$ . We assume that the initial data in (1.3) satisfying  $\mathbf{v}^0 = (v_1^0, v_2^0, v_3^0, v_4^0, v_5^0) := (R^0, S^0, u_1^0, u_2^0, Z^0) \in \mathbb{X}$ . For the local existence and positivity of solutions in the space  $\mathbb{X}$ , we appeal to the theory developed in [8] where existence and uniqueness and positivity are treated simultaneously (taking delay as zero). The idea is to view the system (1.1)–(1.3) as the abstract ordinary differential equation in  $\mathbb{X}$  and the so-called mild solutions can be obtained for any given initial data. More precisely,

$$\begin{cases} v_1(t) = V_R(t, 0)v_1^0 + \int_0^t T(t-s)B_1(\mathbf{v}(s))ds, \\ v_2(t) = V_S(t, 0)v_2^0 + \int_0^t T(t-s)B_2(\mathbf{v}(s))ds, \\ v_i(t) = T(t)v_i^0 + \int_0^t T(t-s)B_i(\mathbf{v}(s))ds, \quad i = 3, 4, 5, \end{cases}$$

where  $T(t)$  is the positive, non-expansive, analytic semigroup on  $C([0, 1], \mathbb{R})$  (see, e.g., [13, Chapter 7]) such that  $v = T(t)v^0$  satisfies the linear initial value problem

$$\begin{cases} \frac{\partial v}{\partial t} = d \frac{\partial^2 v}{\partial x^2}, & t > 0, 0 < x < 1, \\ -\frac{\partial v}{\partial x}(0, t) = 0, \quad \frac{\partial v}{\partial x}(1, t) + \gamma v(1, t) = 0, & t > 0, \\ v(x, 0) = v^0(x). \end{cases}$$

For  $N = R, S$ , we assume that  $V_N(t, s), t > s$ , is the family of affine operators on  $C([0, 1], \mathbb{R})$  (see, e.g., [11, Chapter 5]) such that  $v = V_N(t, s)v^0$  satisfies the linear system with nonhomogeneous boundary conditions, with start time  $s$ , given by

$$\begin{cases} \frac{\partial v}{\partial t} = d \frac{\partial^2 v}{\partial x^2}, & t > 0, 0 < x < 1, \\ -\frac{\partial v}{\partial x}(0, t) = N^{(0)}, \quad \frac{\partial v}{\partial x}(1, t) + \gamma v(1, t) = 0, & t > s, \\ v(x, s) = v^0(x). \end{cases}$$

The nonlinear operators  $B_1, B_2, B_i : C([0, 1], \mathbb{R}_+) \rightarrow C([0, 1], \mathbb{R}), i = 3, 4, 5$ , are defined by

$$\begin{cases} B_1(\mathbf{v}) = -q_{1r}f_1(v_1, v_2)v_3e^{-\alpha v_4} - q_{2r}f_2(v_1, v_2)v_4, \\ B_2(\mathbf{v}) = -q_{1s}f_1(v_1, v_2)v_3e^{-\alpha v_4} - q_{2s}f_2(v_1, v_2)v_4, \\ B_3(\mathbf{v}) = f_1(v_1, v_2)v_3e^{-\alpha v_4} - q_1g_1(v_3)v_5, \\ B_4(\mathbf{v}) = f_2(v_1, v_2)v_4 - q_2g_2(v_4)v_5, \\ B_5(\mathbf{v}) = G(v_3, v_4)v_5. \end{cases}$$

By standard maximum principle arguments (see, e.g., [13, Chapter 7]), it follows that  $V_N(t, s)C([0, 1], \mathbb{R}_+) \subset C([0, 1], \mathbb{R}_+), \forall t > s, N = R, S$  and  $T(t)C([0, 1], \mathbb{R}_+) \subset C([0, 1], \mathbb{R}_+), \forall t > 0$ . Since  $g_1(0) = 0$  and  $g_2(0) = 0$ , it follows that  $B_i(\mathbf{v}) \geq 0$  whenever  $v_i \equiv 0, i = 1, 2, 3, 4, 5$ . Hence,  $\mathbf{B} := (B_1, B_2, B_3, B_4, B_5)$  is quasipositive (see, e.g., [8, Remark 1.1]). By [8, Theorem 1 and Remark 1.1], it follows that the system (1.1)–(1.3) has a unique noncontinuable solution and the solutions to (1.1)–(1.3) remain non-negative on their interval of existence if they are non-negative initially. More precisely, we have the following results:

**Lemma 2.1.** *For every initial value function  $\mathbf{v}^0 \in \mathbb{X} = C([0, 1], \mathbb{R}_+^5)$ , system (1.1)–(1.3) has a unique mild solution  $\mathbf{v}(x, t, \mathbf{v}^0)$  on  $(0, \tau_{\mathbf{v}^0})$  with  $\mathbf{v}(\cdot, 0, \mathbf{v}^0) = \mathbf{v}^0$ , where  $\tau_{\mathbf{v}^0} \leq \infty$ . Furthermore,  $\mathbf{v}(\cdot, t, \mathbf{v}^0) \in \mathbb{X}, \forall t \in (0, \tau_{\mathbf{v}^0})$  and  $\mathbf{v}(x, t, \mathbf{v}^0)$  is a classical solution of (1.1)–(1.3),  $\forall t > 0$ .*

For  $N = R, S$ , we consider

$$\begin{cases} \frac{\partial W_N}{\partial t} = d \frac{\partial^2 W_N}{\partial x^2}, & x \in (0, 1), t > 0, \\ \frac{\partial W_N}{\partial x}(0, t) = -N^{(0)}, \quad \frac{\partial W_N}{\partial x}(1, t) + \gamma W_N(1, t) = 0, & t > 0, \\ W_N(x, 0) = W_N^0(x), & x \in (0, 1). \end{cases} \tag{2.1}$$

Then  $W_N(x, t)$  satisfies (see, e.g., [3])

$$\lim_{t \rightarrow \infty} W_N(x, t) = w_N(x), \text{ uniformly in } x \in [0, 1], \tag{2.2}$$

where  $w_N(x) = N^{(0)}(\frac{1+\gamma}{\gamma} - x)$ .

Next, we show that solutions of (1.1)–(1.3) are ultimately bounded.

**Lemma 2.2.** Any solution with initial value function in  $\mathbb{X}$  of the system (1.1)–(1.3) exists globally on  $[0, \infty)$ . Moreover, solutions are ultimately bounded.

**Proof.** We first consider the case where  $G(u_1, u_2)$  takes the form (1.5). Let

$$Y_R(x, t) = R(x, t) + q_{1r}u_1(x, t) + q_{2r}u_2(x, t) + \min\{q_1q_{1r}, q_2q_{2r}\}Z,$$

and

$$Y_S(x, t) = S(x, t) + q_{1s}u_1(x, t) + q_{2s}u_2(x, t) + \min\{q_1q_{1s}, q_2q_{2s}\}Z.$$

Then  $Y_R(x, t)$  satisfies

$$\begin{aligned} \frac{\partial Y_R}{\partial t} &= d \frac{\partial^2 Y_R}{\partial x^2} + \{\min\{q_1q_{1r}, q_2q_{2r}\} \min\{g_1(u_1), g_2(u_2)\} - q_1q_{1r}g_1(u_1) - q_2q_{2r}g_2(u_2)\}Z \\ &\leq d \frac{\partial^2 Y_R}{\partial x^2}, \quad x \in (0, 1), \quad t > 0. \end{aligned}$$

That is,  $Y_R(x, t)$  satisfies

$$\begin{cases} \frac{\partial Y_R}{\partial t} \leq d \frac{\partial^2 Y_R}{\partial x^2}, \quad x \in (0, 1), \quad t > 0, \\ \frac{\partial Y_R}{\partial x}(0, t) = -R^{(0)}, \quad \frac{\partial Y_R}{\partial x}(1, t) + \gamma Y_R(1, t) = 0, \quad t > 0, \\ Y_R(x, 0) = Y_R^0(x), \quad x \in (0, 1). \end{cases} \tag{2.3}$$

Comparing system (2.3) with (2.1), it follows that  $Y_R(x, t) \leq W_R(x, t)$ ,  $x \in [0, 1]$ ,  $t > 0$ , where we have put  $N = R$  and  $Y_R^0(\cdot) \equiv W_R^0(\cdot)$  in (2.1). Thus,

$$\limsup_{t \rightarrow \infty} Y_R(x, t) \leq w_R(x) := R^{(0)}\left(\frac{1 + \gamma}{\gamma} - x\right), \quad x \in [0, 1]. \tag{2.4}$$

Similarly, we can show that

$$\limsup_{t \rightarrow \infty} Y_S(x, t) \leq w_S(x) := S^{(0)}\left(\frac{1 + \gamma}{\gamma} - x\right), \quad x \in [0, 1]. \tag{2.5}$$

From Lemma 2.1, (2.4), and (2.5), it follows that  $R, S, u_1, u_2$ , and  $Z$  are ultimately bounded.

For the case where  $G(u_1, u_2)$  takes the form (1.4), it follows that

$$G(u_1, u_2) = g_1(u_1) \cdot g_2(u_2) \leq m_{1z}g_2(u_2).$$

Setting

$$Y_R(x, t) = R(x, t) + q_{1r}u_1(x, t) + q_{2r}u_2(x, t) + \frac{q_2q_{2r}}{m_{1z}}Z,$$

and

$$Y_S(x, t) = S(x, t) + q_{1s}u_1(x, t) + q_{2s}u_2(x, t) + \frac{q_2q_{2s}}{m_{1z}}Z.$$

By the same arguments as before, we are able to show that  $Y_R(x, t)$  and  $Y_S(x, t)$  are ultimately bounded. So are  $R, S, u_1, u_2$ , and  $Z$ .  $\square$

By the strong maximum principle and the Hopf boundary lemma (see [12]), we have the following result.

**Lemma 2.3.** *Let*

$$(v_1(x, t), v_2(x, t), v_3(x, t), v_4(x, t), v_5(x, t)) := (R(x, t), S(x, t), u_1(x, t), u_2(x, t), Z(x, t))$$

be the solution of system (1.1)–(1.3) with initial data  $\mathbf{v}^0 \in \mathbb{X}$ . If there is a  $t_0 \geq 0$  such that  $v_i(\cdot, t_0) \not\equiv 0$ , for some  $i \in \{1, 2, 3, 4, 5\}$ , then  $v_i(x, t) > 0$ , for all  $x \in [0, 1]$  and  $t > t_0$ .

### 3. Dynamics of harmful algae

In this section, we put  $Z = 0$  in (1.1)–(1.3) and consider the following system:

$$\begin{cases} \frac{\partial R}{\partial t} = d \frac{\partial^2 R}{\partial x^2} - q_{1r} f_1(R, S) u_1 e^{-\alpha u_2} - q_{2r} f_2(R, S) u_2, & x \in (0, 1), t > 0, \\ \frac{\partial S}{\partial t} = d \frac{\partial^2 S}{\partial x^2} - q_{1s} f_1(R, S) u_1 e^{-\alpha u_2} - q_{2s} f_2(R, S) u_2, & x \in (0, 1), t > 0, \\ \frac{\partial u_1}{\partial t} = d \frac{\partial^2 u_1}{\partial x^2} + f_1(R, S) u_1 e^{-\alpha u_2}, & x \in (0, 1), t > 0, \\ \frac{\partial u_2}{\partial t} = d \frac{\partial^2 u_2}{\partial x^2} + f_2(R, S) u_2, & x \in (0, 1), t > 0, \end{cases} \tag{3.1}$$

with boundary conditions

$$\begin{cases} \frac{\partial R}{\partial x}(0, t) = -R(0), \quad \frac{\partial R}{\partial x}(1, t) + \gamma R(1, t) = 0, & t > 0, \\ \frac{\partial S}{\partial x}(0, t) = -S(0), \quad \frac{\partial S}{\partial x}(1, t) + \gamma S(1, t) = 0, & t > 0, \\ \frac{\partial u_i}{\partial x}(0, t) = \frac{\partial u_i}{\partial x}(1, t) + \gamma u_i(1, t) = 0, & t > 0, \quad i = 1, 2, \end{cases} \tag{3.2}$$

and initial conditions

$$\begin{cases} R(x, 0) = R^0(x) \geq 0, S(x, 0) = S^0(x) \geq 0, & x \in (0, 1), \\ u_i(x, 0) = u_i^0(x) \geq 0, & x \in (0, 1), \quad i = 1, 2. \end{cases} \tag{3.3}$$

#### 3.1. A single population model

For  $i = 1, 2$ , we first consider the following system related to the single population model of (3.1)–(3.3):

$$\begin{cases} \frac{\partial R}{\partial t} = d \frac{\partial^2 R}{\partial x^2} - q_{ir} f_i(R, S) u_i, & x \in (0, 1), t > 0, \\ \frac{\partial S}{\partial t} = d \frac{\partial^2 S}{\partial x^2} - q_{is} f_i(R, S) u_i, & x \in (0, 1), t > 0, \\ \frac{\partial u_i}{\partial t} = d \frac{\partial^2 u_i}{\partial x^2} + f_i(R, S) u_i, & x \in (0, 1), t > 0, \end{cases} \tag{3.4}$$

with boundary conditions

$$\begin{cases} \frac{\partial R}{\partial x}(0, t) = -R(0), \quad \frac{\partial R}{\partial x}(1, t) + \gamma R(1, t) = 0, & t > 0, \\ \frac{\partial S}{\partial x}(0, t) = -S(0), \quad \frac{\partial S}{\partial x}(1, t) + \gamma S(1, t) = 0, & t > 0, \\ \frac{\partial u_i}{\partial x}(0, t) = \frac{\partial u_i}{\partial x}(1, t) + \gamma u_i(1, t) = 0, & t > 0, \end{cases} \tag{3.5}$$

and initial conditions

$$\begin{cases} R(x, 0) = R^0(x) \geq 0, S(x, 0) = S^0(x) \geq 0, & x \in (0, 1), \\ u_i(x, 0) = u_i^0(x) \geq 0, & x \in (0, 1). \end{cases} \tag{3.6}$$

Note that if we put  $u_2 = 0$  ( $u_1 = 0$ ) in (3.1)–(3.3), then we obtain the system (3.4)–(3.6) with  $i = 1$  ( $i = 2$ ).

For  $i = 1, 2$ , introducing the new variable  $W_{iR}(x, t) = R(x, t) + q_{ir}u_i(x, t)$  into (3.4)–(3.6), then  $W_{iR}(x, t)$  satisfies (2.1) with  $N = R$ . Hence,

$$\lim_{t \rightarrow \infty} W_{iR}(x, t) = w_R(x) := R^{(0)}\left(\frac{1 + \gamma}{\gamma} - x\right), \text{ uniformly in } x \in [0, 1].$$

Similarly, introducing the new variable  $W_{iS}(x, t) = S(x, t) + q_{is}u_i(x, t)$  into (3.4)–(3.6), it follows that

$$\lim_{t \rightarrow \infty} W_{iS}(x, t) = w_S(x) := S^{(0)}\left(\frac{1 + \gamma}{\gamma} - x\right), \text{ uniformly in } x \in [0, 1].$$

Then we conclude the limiting system of (3.4)–(3.6) takes the form,  $i = 1, 2$ ,

$$\begin{cases} \frac{\partial u_i}{\partial t} = d \frac{\partial^2 u_i}{\partial x^2} + f_i(w_R(x) - q_{ir}u_i, w_S(x) - q_{is}u_i)u_i, & x \in (0, 1), t > 0, \\ \frac{\partial u_i}{\partial x}(0, t) = \frac{\partial u_i}{\partial x}(1, t) + \gamma u_i(1, t) = 0, & t > 0, \\ u_i(x, 0) = u_i^0(x) \geq 0, & x \in (0, 1). \end{cases} \tag{3.7}$$

We denote  $\lambda_i^0$  to be the principal eigenvalue of the eigenvalue problem

$$\begin{cases} d\varphi''(x) + f_i(w_R(x), w_S(x))\varphi(x) = \lambda_i\varphi(x), & x \in (0, 1), \\ \varphi'(0) = \varphi'(1) + \gamma\varphi(1) = 0, \end{cases} \tag{3.8}$$

with the corresponding positive eigenfunctions uniquely determined by the normalization.

By similar arguments to those in [10,16,18], we have the following results on the global stability of system (3.7).

**Lemma 3.1.** *For any nonnegative initial function  $u_i^0(x)$ ,  $i = 1, 2$ , with  $q_{ir}u_i^0(x) \leq w_R(x)$  and  $q_{is}u_i^0(x) \leq w_S(x)$ , there exists a unique nonnegative solution  $u_i(x, t)$  of (3.7) defined for  $t > 0$ . Furthermore,*

- (i) *If  $\lambda_i^0 \leq 0$ , then  $\lim_{t \rightarrow \infty} u_i(x, t) = 0$ , uniformly for  $x \in [0, 1]$ .*
- (ii) *If  $\lambda_i^0 > 0$ , then there exists a unique positive solution  $u_i^*(x)$  with  $q_{ir}u_i^*(x) < w_R(x)$  and  $q_{is}u_i^*(x) < w_S(x)$  on  $[0, 1]$  such that*

$$\lim_{t \rightarrow \infty} u_i(x, t) = u_i^*(x),$$

*uniformly for  $x \in [0, 1]$  provided that  $u_i^0(\cdot) \not\equiv 0$ .*

With Lemma 3.1, we can adopt the arguments in [17, Theorem 2.2] to lift the dynamics of the limiting system (3.7) to the system (3.4)–(3.6) by using the theory of chain transitive sets [15,19].

**Lemma 3.2.** *For any nonnegative initial function  $(R^0(x), S^0(x), u_i^0(x))$  with  $u_i^0(x) \geq 0$ , there exists a unique nonnegative solution  $(R(x, t), S(x, t), u_i(x, t))$  of (3.4)–(3.6) defined for  $t > 0$ . Furthermore,*

- (i) *If  $\lambda_i^0 \leq 0$ , then  $\lim_{t \rightarrow \infty} (R(x, t), S(x, t), u_i(x, t)) = (w_R(x), w_S(x), 0)$ , uniformly for  $x \in [0, 1]$ .*
- (ii) *If  $\lambda_i^0 > 0$ , then there exists a unique positive solution  $(R_i^*(x), S_i^*(x), u_i^*(x))$  on  $[0, 1]$  such that*

$$\lim_{t \rightarrow \infty} (R(x, t), S(x, t), u_i(x, t)) = (R_i^*(x), S_i^*(x), u_i^*(x)),$$

uniformly for  $x \in [0, 1]$  provided that  $u_i^0(\cdot) \not\equiv 0$ . Here,  $u_i^*(x)$  is defined in Lemma 3.1, and

$$R_i^*(x) = w_R(x) - q_{ir}u_i^*(x), \quad S_i^*(x) = w_S(x) - q_{is}u_i^*(x), \quad i = 1, 2. \quad (3.9)$$

### 3.2. Coexistence of harmful algae

This section is devoted to the study of the global dynamics of system (3.1)–(3.3). From Lemma 3.2, it is easy to see that system (3.1)–(3.3) has the following possible steady-state solutions:

- (i) Trivial steady-state solution  $E_0 = (R, S, u_1, u_2) = (w_R(x), w_S(x), 0, 0)$  always exists;
- (ii) Semi-trivial steady-state solution  $E_1 = (R, S, u_1, u_2) = (R_1^*(x), S_1^*(x), u_1^*(x), 0)$  exists provided that  $\lambda_1^0 > 0$ , where  $(R_1^*(x), S_1^*(x), u_1^*(x))$  is the unique steady-state solution of system (3.4)–(3.6) with  $i = 1$  (see Lemma 3.2 (ii));
- (iii) Semi-trivial steady-state solution  $E_2 = (R, S, u_1, u_2) = (R_2^*(x), S_2^*(x), 0, u_2^*(x))$  exists provided that  $\lambda_2^0 > 0$ , where  $(R_2^*(x), S_2^*(x), u_2^*(x))$  is the unique steady-state solution of system (3.4)–(3.6) with  $i = 2$  (see Lemma 3.2 (ii)).

Of course, there may be additional steady-state solutions as well and these must be positive. The two algae can coexist if a positive steady-state solution exists.

In order to investigate the local stability of  $E_1$  for system (3.1)–(3.3), we consider the following linear system:

$$\begin{cases} \frac{\partial u_2}{\partial t} = d \frac{\partial^2 u_2}{\partial x^2} + f_2(R_1^*(x), S_1^*(x))u_2, & x \in (0, 1), \quad t > 0, \\ \frac{\partial u_2}{\partial x}(0, t) = \frac{\partial u_2}{\partial x}(1, t) + \gamma u_2(1, t) = 0, & t > 0, \\ u_2(x, 0) = u_2^0(x) \geq 0, & x \in (0, 1). \end{cases} \quad (3.10)$$

Then we denote  $\Lambda_1^0$  to be the principal eigenvalue of the eigenvalue problem

$$\begin{cases} d\varphi''(x) + f_2(R_1^*(x), S_1^*(x))\varphi(x) = \Lambda_1\varphi(x), & x \in (0, 1), \\ \varphi'(0) = \varphi'(1) + \gamma\varphi(1) = 0, \end{cases} \quad (3.11)$$

with the corresponding positive eigenfunctions uniquely determined by the normalization. For the local stability of  $E_2$  for system (3.1)–(3.3), we consider the following linear system:

$$\begin{cases} \frac{\partial u_1}{\partial t} = d \frac{\partial^2 u_1}{\partial x^2} + f_1(R_2^*(x), S_2^*(x))e^{-\alpha u_2^*(x)}u_1, & x \in (0, 1), \quad t > 0, \\ \frac{\partial u_1}{\partial x}(0, t) = \frac{\partial u_1}{\partial x}(1, t) + \gamma u_1(1, t) = 0, & t > 0, \\ u_1(x, 0) = u_1^0(x) \geq 0, & x \in (0, 1). \end{cases} \quad (3.12)$$

Then we denote  $\Lambda_2^0$  to be the principal eigenvalue of the eigenvalue problem

$$\begin{cases} d\varphi''(x) + f_1(R_2^*(x), S_2^*(x))e^{-\alpha u_2^*(x)}\varphi(x) = \Lambda_2\varphi(x), & x \in (0, 1), \\ \varphi'(0) = \varphi'(1) + \gamma\varphi(1) = 0, \end{cases} \quad (3.13)$$

with the corresponding positive eigenfunctions uniquely determined by the normalization.

Introducing the new variable  $W_R(x, t) = R(x, t) + q_{1r}u_1(x, t) + q_{2r}u_2(x, t)$ , and  $W_S(x, t) = S(x, t) + q_{1s}u_1(x, t) + q_{2s}u_2(x, t)$  into (3.1)–(3.3), respectively. Then  $W_N(x, t)$  satisfies (2.1) with  $N = R, S$ , and hence,



$$\lim_{t \rightarrow \infty} W_N(x, t) = w_N(x) := N^{(0)}\left(\frac{1 + \gamma}{\gamma} - x\right), \text{ uniformly in } x \in [0, 1]. \tag{3.14}$$

Let  $\Omega = C([0, 1], \mathbb{R}_+^4)$  be the positive cone of the Banach space  $C([0, 1], \mathbb{R}^4)$  with the usual supremum norm. By (3.14) and similar arguments to those in Lemma 2.2, we can show that any solution with initial value function in  $\Omega$  of the system (3.1)–(3.3) exists globally on  $[0, \infty)$ , and solutions are ultimately bounded. Let  $\Pi(t) : \Omega \rightarrow \Omega$  be the semiflow generated by system (3.1)–(3.3). Then  $\Pi(t) : \Omega \rightarrow \Omega$  is compact, point dissipative,  $\forall t > 0$ . By [2, Theorem 3.4.8], it follows that  $\Pi(t)$  admits a global compact attractor in  $\Omega$ . Setting

$$\Omega_0 = \{(R^0(\cdot), S^0(\cdot), u_1^0(\cdot), u_2^0(\cdot)) \in \Omega : u_1^0(\cdot) \not\equiv 0, u_2^0(\cdot) \not\equiv 0\}, \quad \partial\Omega_0 = \Omega \setminus \Omega_0.$$

**Theorem 3.1.** *Assume that*

$$\lambda_i^0 > 0 \text{ and } \Lambda_i^0 > 0, \quad i = 1, 2. \tag{3.15}$$

*Then  $\Pi(t) : \Omega \rightarrow \Omega$  is uniformly persistent with respect to  $(\Omega_0, \partial\Omega_0)$  in the sense that there exists  $\tilde{\eta} > 0$  such that*

$$\liminf_{t \rightarrow \infty} u_i(\cdot, t, Q^0) \geq \tilde{\eta}, \quad i = 1, 2, \quad \forall Q^0 \in \Omega_0.$$

*Further, system (3.1)–(3.3) admits at least one positive steady-state solutions*

$$(\tilde{R}(x), \tilde{S}(x), \tilde{u}_1(x), \tilde{u}_2(x)).$$

**Proof.** By the same arguments in Lemma 2.3, it follows that for any  $Q^0 \in \Omega_0$ , we have

$$u_1(x, t, Q^0) > 0, \quad u_2(x, t, Q^0) > 0, \quad \forall x \in [0, 1], \quad t > 0.$$

This implies that  $\Pi(t)\Omega_0 \subseteq \Omega_0$  for all  $t \geq 0$ . Let

$$\tilde{M}_\partial := \{Q^0 \in \partial\Omega_0 : \Pi(t)Q^0 \in \partial\Omega_0, \forall t \geq 0\},$$

and  $\tilde{\omega}(Q^0)$  be the omega limit set of the orbit  $O^+(Q^0) := \{\Pi(t)Q^0 : t \geq 0\}$ . We further prove the following claims.

*Claim 1.*  $\tilde{\omega}(\varphi) \subset \tilde{M}_0 \cup \tilde{M}_1 \cup \tilde{M}_2, \forall \varphi \in \tilde{M}_\partial$ , where  $\tilde{M}_i = \{E_i\}, i = 0, 1, 2$ .

Since  $\varphi \in \tilde{M}_\partial$ , we have  $\Pi(t)\varphi \in \tilde{M}_\partial, \forall t \geq 0$ . Thus  $u_1(\cdot, t, \varphi) \equiv 0$  or  $u_2(\cdot, t, \varphi) \equiv 0, \forall t \geq 0$ . In case where  $u_1(\cdot, t, \varphi) \equiv 0, \forall t \geq 0$ . Then  $(R, S, u_2)$  satisfies (3.4)–(3.6) with  $i = 2$ . By Lemma 3.2, it follows that either

$$\lim_{t \rightarrow \infty} (R(x, t, \varphi), S(x, t, \varphi), u_2(x, t, \varphi)) = (w_R(x), w_S(x), 0), \text{ uniformly for } x \in [0, 1],$$

or

$$\lim_{t \rightarrow \infty} (R(x, t, \varphi), S(x, t, \varphi), u_2(x, t, \varphi)) = (R_2^*(x), S_2^*(x), u_2^*(x)), \text{ uniformly for } x \in [0, 1].$$

In case where  $u_1(\cdot, t_0, \varphi) \not\equiv 0$ , for some  $t_0 \geq 0$ . Then the strong maximum principle and the Hopf boundary lemma (see [12]) implies that  $u_1(\cdot, t, \varphi) > 0$ , for all  $t > t_0$ . Thus,  $u_2(\cdot, t, \varphi) \equiv 0, \forall t > t_0$ , and hence, the

$(R, S, u_1)$  equation in (3.1)–(3.3) satisfies (3.4)–(3.6) with  $i = 1$ , for all  $t > t_0$ . Again, from Lemma 3.2, it follows that either

$$\lim_{t \rightarrow \infty} (R(x, t, \varphi), S(x, t, \varphi), u_1(x, t, \varphi)) = (w_R(x), w_S(x), 0), \text{ uniformly for } x \in [0, 1],$$

or

$$\lim_{t \rightarrow \infty} (R(x, t, \varphi), S(x, t, \varphi), u_1(x, t, \varphi)) = (R_1^*(x), S_1^*(x), u_1^*(x)), \text{ uniformly for } x \in [0, 1].$$

This ends the proof of Claim 1.

*Claim 2.* For  $i = 0, 1, 2$ ,  $\tilde{M}_i$  is a uniform weak repeller for  $\Omega_0$  in the sense that there exists  $\tilde{\delta}_i > 0$  such that  $\limsup_{t \rightarrow \infty} \|\Pi(t)Q - \tilde{M}_i\| \geq \tilde{\delta}_i, \forall Q \in \Omega_0$ .

We only show the case where  $i = 2$  since the other cases can be proved in a similar way. From the fact that  $\Lambda_2^0 > 0$ , we may assume that there exists an  $\epsilon_0 > 0$  such that  $\Lambda_2^{\epsilon_0} > 0$ , where  $\Lambda_2^{\epsilon_0}$  is the principal eigenvalue of the eigenvalue problem

$$\begin{cases} d\varphi''(x) + [f_1(R_2^*(x), S_2^*(x))e^{-\alpha u_2^*(x)} - \epsilon_0]\varphi(x) = \Lambda_2\varphi(x), & x \in (0, 1), \\ \varphi'(0) = \varphi'(1) + \gamma\varphi(1) = 0. \end{cases}$$

The positive eigenfunction corresponding to  $\Lambda_2^{\epsilon_0}$  can be uniquely determined by the normalization, and we denote it by  $\varphi_{\epsilon_0}(x)$ . By continuity of  $f_1$ , we can choose  $\tilde{\delta}_2 > 0$  such that if  $\|(R_2, S_2, u_2) - (R_2^*(\cdot), S_2^*(\cdot), u_2^*(\cdot))\| < \tilde{\delta}_2$ , then

$$f_1(R_2, S_2)e^{-\alpha u_2} > f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - \epsilon_0. \tag{3.16}$$

Then we show that

$$\limsup_{t \rightarrow \infty} \|\Pi(t)Q - \tilde{M}_2\| \geq \tilde{\delta}_2, \forall Q \in \Omega_0. \tag{3.17}$$

Suppose, by contradiction, there exists  $Q^0 \in \Omega_0$  such that  $\limsup_{t \rightarrow \infty} \|\Pi(t)Q^0 - \tilde{M}_2\| < \tilde{\delta}_2$ . Then there exists  $t_0 > 0$  such that for  $t \geq t_0$  and  $x \in [0, 1]$ , we have

$$\|(R(x, t, Q^0), S(x, t, Q^0), u_2(x, t, Q^0)) - (R_2^*(x), S_2^*(x), u_2^*(x))\| < \tilde{\delta}_2.$$

It then follows from (3.16) that

$$f_1(R(\cdot, t, Q^0), S(\cdot, t, Q^0))e^{-\alpha u_2(\cdot, t, Q^0)} > f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - \epsilon_2, \quad t \geq t_0. \tag{3.18}$$

With (3.18), it follows from the third equation of (1.1) that

$$\begin{cases} \frac{\partial u_1}{\partial t} \geq d\frac{\partial^2 u_1}{\partial x^2} + [f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - \epsilon_0]u_1, & x \in (0, 1), \quad t \geq t_0, \\ \frac{\partial u_1}{\partial x}(0, t) = \frac{\partial u_1}{\partial x}(1, t) + \gamma u_1(1, t) = 0, & t \geq t_0. \end{cases} \tag{3.19}$$

Since  $Q^0 \in \Omega_0$ , we can further show that  $u_1(x, t, Q^0) > 0, \forall x \in [0, 1], t > 0$ . Thus, there exists a sufficiently small number  $a > 0$  such that  $u_1(x, t_0, Q^0) \geq a\varphi_{\epsilon_0}(x), \forall x \in [0, 1]$ . Note that  $\hat{u}_1(x, t) := ae^{\Lambda_2^{\epsilon_0}(t-t_0)}\varphi_{\epsilon_0}(x), t \geq t_0$ , is a solution of the following linear system:

$$\begin{cases} \frac{\partial u_1}{\partial t} = d \frac{\partial^2 u_1}{\partial x^2} + [f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - \epsilon_0]u_1, & x \in (0, 1), t \geq t_0, \\ \frac{\partial u_1}{\partial x}(0, t) = \frac{\partial u_1}{\partial x}(1, t) + \gamma u_1(1, t) = 0, & t \geq t_0, \end{cases} \tag{3.20}$$

with initial data  $\hat{u}_1(x, t_0) := a\varphi_{\epsilon_0}(x)$ . Then the comparison principle implies that

$$u_1(x, t, Q^0) \geq \hat{u}_1(x, t) := ae^{\Lambda_2^{\epsilon_0}(t-t_0)}\varphi_{\epsilon_0}(x), \quad \forall x \in [0, 1], t \geq t_0.$$

Since  $\Lambda_2^{\epsilon_0} > 0$ , it follows that  $u_1(x, t, Q^0)$  is unbounded. This contradiction proves the result in (3.17). Thus, Claim 2 holds.

Define a continuous function  $\rho : \Omega \rightarrow [0, \infty)$  by

$$\rho(Q) := \min_{3 \leq i \leq 4} \{ \min_{x \in [0,1]} Q_i(x) \}, \quad \forall Q := (Q_1, Q_2, Q_3, Q_4) \in \Omega.$$

By the strong maximum principle and the Hopf boundary lemma (see [12]), we can prove that  $\rho^{-1}(0, \infty) \subseteq \Omega_0$  and  $\rho$  has the property that if  $\rho(Q) > 0$  or  $Q \in \Omega_0$  with  $\rho(Q) = 0$ , then  $\rho(\Pi(t)Q) > 0, \forall t > 0$ . That is,  $\rho$  is a generalized distance function for the semiflow  $\Pi(t) : \Omega \rightarrow \Omega$  (see, e.g., [15]). By the above claims, it follows that any forward orbit of  $\Pi(t)$  in  $\tilde{M}_\partial$  converges to either  $\tilde{M}_0$  or  $\tilde{M}_1$  or  $\tilde{M}_2$ . Further,  $\tilde{M}_0, \tilde{M}_1$ , and  $\tilde{M}_2$  are isolated in  $\Omega$  and  $W^s(\tilde{M}_i) \cap \Omega_0 = \emptyset, \forall i = 0, 1, 2$ , where  $W^s(\tilde{M}_i)$  is the stable set of  $\tilde{M}_i, i = 0, 1, 2$  (see [15]). It is easy that no subsets of  $\tilde{M}_0, \tilde{M}_1, \tilde{M}_2$  forms a cycle in  $\tilde{M}_\partial$ .

Since  $\Pi(t) : \Omega \rightarrow \Omega$  admits a global compact attractor in  $\Omega$ , it follows from [15, Theorem 3] that there exists an  $\tilde{\eta} > 0$  such that

$$\min_{\varphi \in \omega(Q)} \rho(\varphi) > \tilde{\eta}, \quad \forall Q \in \Omega_0.$$

This implies that the uniform persistence stated in our theorem is valid. By [9, Theorem 3.7 and Remark 3.10], it then follows that  $\Pi(t) : \Omega_0 \rightarrow \Omega_0$  has a global attractor. It then follows from [9, Theorem 4.7] that  $\Pi(t)$  has an steady-state solution  $(\hat{R}(\cdot), \hat{S}(\cdot), \hat{u}_1(\cdot), \hat{u}_2(\cdot), \hat{Z}(\cdot)) \in \Omega_0. \quad \square$

**Remark 3.1.** From Theorem 3.1, we see that  $\Pi(t) : \Omega \rightarrow \Omega$  is uniformly persistent with respect to  $(\Omega_0, \partial\Omega_0)$  provided that (3.15) is met. It follows from [9, Theorem 3.8] that  $\Pi(t) : \Omega_0 \rightarrow \Omega_0$  admits a global attractor  $A_0$ . Since  $A_0 \subset \Omega_0$  and  $A_0 = \Pi(t)(A_0)$ , we further have  $A_0 \subset \text{Int}(C([0, 1], \mathbb{R}_+^4))$ .

#### 4. Coexistence of harmful algae and zooplankton

In this section, we explore the possibility of coexistence of harmful algae and zooplankton. That is, we are going to establish the existence of positive (coexistence) steady-state solutions of system (1.1)–(1.3). Since zooplankton population growth rate  $G(u_1, u_2)$  takes the form (1.4) or (1.5), it follows that  $G(0, u_2) = G(u_1, 0) = 0$ . This implies that the following types of steady-state solutions of (1.1)–(1.3) cannot occur:

$$(R, S, u_1, u_2, Z) = (\check{R}_1(x), \check{S}_1(x), \check{u}_1(x)(x), 0, \check{Z}_1(x)) \text{ with } \check{u}_1(\cdot), \check{Z}_1(\cdot) \gg 0, \tag{4.1}$$

and

$$(R, S, u_1, u_2, Z) = (\check{R}_2(x), \check{S}_2(x), 0, \check{u}_2(x)(x), \check{Z}_2(x)) \text{ with } \check{u}_2(\cdot), \check{Z}_2(\cdot) \gg 0. \tag{4.2}$$

Thus, system (1.1)–(1.3) has the following possible steady-state solutions:

- (i) Trivial steady-state solution  $\mathcal{E}_0 = (R, S, u_1, u_2, Z) = (w_R(x), w_S(x), 0, 0, 0)$  always exists (see Lemma 3.2 (i));

- (ii) Semi-trivial steady-state solution  $\mathcal{E}_1 = (R, S, u_1, u_2, Z) = (R_1^*(x), S_1^*(x), u_1^*(x), 0, 0)$  exists provided that  $\lambda_1^0 > 0$ , where  $(R_1^*(x), S_1^*(x), u_1^*(x))$  is the unique steady-state solution of system (3.4)–(3.6) with  $i = 1$  (see Lemma 3.2 (ii));
- (iii) Semi-trivial steady-state solution  $\mathcal{E}_2 = (R, S, u_1, u_2, Z) = (R_2^*(x), S_2^*(x), 0, u_2^*(x), 0)$  exists provided that  $\lambda_2^0 > 0$ , where  $(R_2^*(x), S_2^*(x), u_2^*(x))$  is the unique steady-state solution of system (3.4)–(3.6) with  $i = 2$  (see Lemma 3.2 (ii));
- (iv) Zooplankton-extinct steady-state solution

$$\mathcal{E}_3 = (R, S, u_1, u_2, Z) = (\tilde{R}(x), \tilde{S}(x), \tilde{u}_1(x), \tilde{u}_2(x), 0)$$

exists provided that (3.15) holds. Here  $(\tilde{R}(x), \tilde{S}(x), \tilde{u}_1(x), \tilde{u}_2(x))$  is a positive steady-state solution of (3.1)–(3.3), which is not necessarily unique (see Theorem 3.1 and Remark 3.1).

Biologically, the most interesting question is whether both harmful algae and zooplankton can coexist in the unstirred chemostat. Mathematically, we want to show the existence of positive (coexistence) steady-state solutions of system (1.1)–(1.3) under suitable conditions.

Recall that  $\mathbb{X} = C([0, 1], \mathbb{R}_+^5)$  is a positive cone of the Banach space  $C([0, 1], \mathbb{R}^5)$  with the usual supremum norm. In view of Lemma 2.1 and Lemma 2.2, we can assume that  $\Sigma(t) : \mathbb{X} \rightarrow \mathbb{X}$  is the semiflow generated by system (1.1)–(1.3). Let

$$\mathbb{X}_0 = \{(R^0(\cdot), S^0(\cdot), u_1^0(\cdot), u_2^0(\cdot), Z^0(\cdot)) \in \mathbb{X} : u_1^0(\cdot) \not\equiv 0, u_2^0(\cdot) \not\equiv 0, Z^0(\cdot) \not\equiv 0\},$$

and

$$\partial\mathbb{X}_0 = \mathbb{X} \setminus \mathbb{X}_0.$$

Assume that  $M_0 = \{\mathcal{E}_0\}$ ,  $M_1 = \{\mathcal{E}_1\}$ ,  $M_2 = \{\mathcal{E}_2\}$ , and  $M_3 = A_0 \times \{0\}$ , where  $A_0 \subset \text{Int}(C([0, 1], \mathbb{R}_+^4))$  is a global attractor of the semiflows generated by system (3.1)–(3.3) (see Remark 3.1).

Define a projection  $\mathcal{P}$  on  $C([0, 1], \mathbb{R}_+^4)$  by

$$\mathcal{P}(R, S, u_1, u_2) = (u_1, u_2), \quad \forall (R, S, u_1, u_2) \in C([0, 1], \mathbb{R}_+^4).$$

Let

$$B_0 = \mathcal{P}(A_0) \text{ and } m(x) = \inf_{\phi \in B_0} G(\phi(x)), \quad \forall x \in [0, 1]. \tag{4.3}$$

**Lemma 4.1.** *Let  $m(x)$  be defined in (4.3). Then  $m(x)$  is continuous on  $[0, 1]$ .*

**Proof.** Let  $H : [0, 1] \times B_0 \rightarrow \mathbb{R}$  be defined by

$$H(x, \phi) = G(\phi(x)), \quad \forall (x, \phi) \in [0, 1] \times B_0.$$

Since  $H$  is continuous on the compact set  $[0, 1] \times B_0$ , it follows that  $H$  is uniformly continuous on  $[0, 1] \times B_0$ . Let  $\epsilon > 0$  to be given. Then there exists  $\delta = \delta(\epsilon)$  such that

$$|G(\phi(x)) - G(\phi(y))| = |H(x, \phi) - H(y, \phi)| < \frac{\epsilon}{2}, \tag{4.4}$$

whenever  $(x, \phi), (y, \phi) \in [0, 1] \times B_0$  with  $|x - y| < \delta$ .

Let  $x, y \in [0, 1]$  be given such that  $|x - y| < \delta$ . Since  $m(y) + \frac{\epsilon}{2}$  is not a lower bound of  $\{G(\phi(y)) : \phi \in B_0\}$ , we can find  $\tilde{\phi} \in B_0$  such that  $m(y) + \frac{\epsilon}{2} > G(\tilde{\phi}(y))$ . Using (4.4), we further have

$$m(y) + \frac{\epsilon}{2} > G(\tilde{\phi}(y)) > G(\tilde{\phi}(x)) - \frac{\epsilon}{2} \geq m(x) - \frac{\epsilon}{2}. \tag{4.5}$$

Similarly,  $m(x) + \frac{\epsilon}{2}$  is not a lower bound of  $\{G(\phi(x)) : \phi \in B_0\}$ , and we can find  $\hat{\phi} \in B_0$  such that  $m(x) + \frac{\epsilon}{2} > G(\hat{\phi}(x))$ . Using (4.4) again, we obtain

$$m(x) + \frac{\epsilon}{2} > G(\hat{\phi}(x)) > G(\hat{\phi}(y)) - \frac{\epsilon}{2} \geq m(y) - \frac{\epsilon}{2}. \tag{4.6}$$

By (4.5) and (4.6), it follows that for any given  $\epsilon > 0$ , if  $x, y \in [0, 1]$  with  $|x - y| < \delta$ , then  $|m(x) - m(y)| < \epsilon$ . This shows that  $m(\cdot)$  is uniformly continuous on  $[0, 1]$ , and hence,  $m(\cdot)$  is continuous on  $[0, 1]$ .  $\square$

Next, we denote  $\mu^0$  to be the principal eigenvalue of the eigenvalue problem

$$\begin{cases} d\psi''(x) + m(x)\psi(x) = \mu\psi(x), & x \in (0, 1), \\ \psi'(0) = \psi'(1) + \gamma\psi(1) = 0 \end{cases} \tag{4.7}$$

where  $m(x)$  is defined in (4.3).

**Lemma 4.2.** *Let (3.15) hold and  $\mu^0 > 0$ . Then  $M_3$  is a uniform weak repeller in the sense that there exists  $\delta_3 > 0$  such that*

$$\limsup_{t \rightarrow \infty} \text{dist}(\Sigma(t)(Q^0, Z^0), M_3) \geq \delta_3, \text{ for all } (Q^0, Z^0) \in \mathbb{X}_0. \tag{4.8}$$

**Proof.** Since  $\mu^0 > 0$ , we can choose a sufficiently small  $\epsilon_0 > 0$  such that  $\mu_{\epsilon_0}^0 > 0$ , where  $\mu_{\epsilon_0}^0$  is the principal eigenvalue of the eigenvalue problem

$$\begin{cases} d\psi''(x) + [m(x) - \epsilon_0]\psi(x) = \mu\psi(x), & x \in (0, 1), \\ \psi'(0) = \psi'(1) + \gamma\psi(1) = 0. \end{cases} \tag{4.9}$$

Define  $\tilde{G} : B_0 \rightarrow C([0, 1], \mathbb{R})$  by

$$\tilde{G}(\phi)(x) = G(\phi(x)), \quad \forall x \in [0, 1], \phi \in B_0.$$

Then there exists  $\delta_3 > 0$  such that

$$\text{dist}(\tilde{G}(\phi), \tilde{G}(B_0)) < \epsilon_0,$$

whenever  $\phi \in C([0, 1], \mathbb{R}^2)$  with  $\text{dist}(\phi, B_0) < \delta_3$ . Since  $B_0$  is compact, it follows that for any  $\phi \in C([0, 1], \mathbb{R}^2)$  with  $\text{dist}(\phi, B_0) < \delta_3$ , there exists  $\phi_* \in B_0$  with  $\phi_*$  depending on  $\phi$  such that

$$\text{dist}(\tilde{G}(\phi), \tilde{G}(\phi_*)) = \text{dist}(\tilde{G}(\phi), \tilde{G}(B_0)) < \epsilon_0.$$

Thus, we have

$$|G(\phi(x)) - G(\phi_*(x))| = |\tilde{G}(\phi)(x) - \tilde{G}(\phi_*)(x)| < \epsilon_0, \quad \forall x \in [0, 1], \tag{4.10}$$

whenever  $\phi \in C([0, 1], \mathbb{R}^2)$  with  $\text{dist}(\phi, B_0) < \delta_3$ .

Next, we prove (4.8) by contradiction. Suppose that (4.8) is not true. Then there exists  $(Q^0, Z^0) \in \mathbb{X}_0$  such that

$$\limsup_{t \rightarrow \infty} \text{dist}(\Sigma(t)(Q^0, Z^0), M_3) < \delta_3,$$

and hence,

$$\limsup_{t \rightarrow \infty} \text{dist}((u_1(\cdot, t), u_2(\cdot, t)), B_0) < \delta_3, \tag{4.11}$$

and

$$\limsup_{t \rightarrow \infty} \|Z(\cdot, t)\| < \delta_3. \tag{4.12}$$

From (4.11), we can choose  $t_0 > 0$  such that

$$\text{dist}((u_1(\cdot, t), u_2(\cdot, t)), B_0) < \delta_3, \quad t \geq t_0. \tag{4.13}$$

By (4.10), it follows that there exists  $\phi_*^t \in B_0$  such that

$$|G((u_1(x, t), u_2(x, t)) - G(\phi_*^t(x)))| < \epsilon_0, \quad \forall x \in [0, 1], \quad t \geq t_0, \tag{4.14}$$

and hence,

$$G((u_1(x, t), u_2(x, t)) > G(\phi_*^t(x)) - \epsilon_0 \geq m(x) - \epsilon_0, \quad \forall x \in [0, 1], \quad t \geq t_0. \tag{4.15}$$

It follows from the fifth equation of (1.1) that

$$\begin{cases} \frac{\partial Z}{\partial t} \geq d \frac{\partial^2 Z}{\partial x^2} + [m(x) - \epsilon_0]Z, & x \in (0, 1), \quad t \geq t_0, \\ \frac{\partial Z}{\partial x}(0, t) = \frac{\partial Z}{\partial x}(1, t) + \gamma Z(1, t) = 0, & t \geq t_0. \end{cases} \tag{4.16}$$

Since  $Z(\cdot, t) \not\equiv 0$ , we can further show that  $Z(\cdot, t_0) \gg 0$ , and hence, there exists a sufficiently small number  $a > 0$  such that  $Z(x, t_0) \geq a\psi_{\epsilon_0}(x)$ ,  $\forall x \in [0, 1]$ , where  $\psi_{\epsilon_0}(x)$  is the eigenfunction corresponding to  $\mu_{\epsilon_0}^0$ . Then the comparison principle ensures that

$$Z(x, t) \geq ae^{\mu_{\epsilon_0}^0(t-t_0)}\psi_{\epsilon_0}(x), \quad \forall x \in [0, 1], \quad t \geq t_0.$$

Since  $\mu_{\epsilon_0}^0 > 0$ , we deduce that  $\lim_{t \rightarrow \infty} Z(\cdot, t) = \infty$ , which contradicts (4.12). This proves (4.8).  $\square$

Now we are in a position to prove the main result of this section.

**Theorem 4.1.** *Let (3.15) hold and  $\mu^0 > 0$ . Then system (1.1)–(1.3) is uniformly persistent with respect to  $(\mathbb{X}_0, \partial\mathbb{X}_0)$  in the following sense that there is a constant  $\eta > 0$  such that every solution  $(R(\cdot, t), S(\cdot, t), u_1(\cdot, t), u_2(\cdot, t), Z(\cdot, t))$  of (1.1)–(1.3) with  $(R(\cdot, 0), S(\cdot, 0), u_1(\cdot, 0), u_2(\cdot, 0), Z(\cdot, 0)) \in \mathbb{X}_0$  satisfying*

$$\liminf_{t \rightarrow \infty} u_i(\cdot, t) \geq \eta, \quad \text{and} \quad \liminf_{t \rightarrow \infty} Z(\cdot, t) \geq \eta, \quad i = 1, 2. \tag{4.17}$$

Furthermore, system (1.1)–(1.3) admits at least one (componentwise) positive steady-state solution  $(\hat{R}(\cdot), \hat{S}(\cdot), \hat{u}_1(\cdot), \hat{u}_2(\cdot), \hat{Z}(\cdot))$ .

**Proof.** By Lemma 2.3, it follows that for any  $\mathbf{v}^0 \in \mathbb{X}_0$ , we have

$$u_1(x, t, \mathbf{v}^0) > 0, \quad u_2(x, t, \mathbf{v}^0) > 0, \quad Z(x, t, \mathbf{v}^0) > 0, \quad \forall x \in [0, 1], \quad t > 0.$$

This implies that  $\Sigma(t)\mathbb{X}_0 \subseteq \mathbb{X}_0$  for all  $t \geq 0$ .

Let

$$M_\partial := \{\mathbf{v}^0 \in \partial\mathbb{X}_0 : \Sigma(t)\mathbf{v}^0 \in \partial\mathbb{X}_0, \forall t \geq 0\},$$

and  $\omega(\mathbf{v}^0)$  be the omega limit set of the orbit  $O^+(\mathbf{v}^0) := \{\Sigma(t)\mathbf{v}^0 : t \geq 0\}$ . We further prove the following claims.

*Claim 1.*  $\omega(\psi) \subset M_0 \cup M_1 \cup M_2 \cup M_3, \forall \psi \in M_\partial$ .

For any given  $\psi := (R^0, S^0, u_1^0, u_2^0, Z^0) \in M_\partial$ , we have  $\Sigma(t)\psi \in M_\partial, \forall t \geq 0$ . Thus, for any given  $t \geq 0$ , we have  $u_1(\cdot, t, \psi) \equiv 0$  or  $u_2(\cdot, t, \psi) \equiv 0$  or  $Z(\cdot, t, \psi) \equiv 0$ .

In the case where  $Z(\cdot, t, \psi) \equiv 0$  for all  $t \geq 0$ , substituting  $Z(\cdot, t, \psi) \equiv 0$  into system (1.1)–(1.3). Then the equations for  $(R, S, u_1, u_2)$  satisfy system (3.1)–(3.3). We discuss the following four subcases:

- (i) If  $u_1^0 \equiv 0, u_2^0 \equiv 0$ , then we have  $u_1(\cdot, t, \psi) \equiv 0$  and  $u_2(\cdot, t, \psi) \equiv 0$ . Thus,  $\lim_{t \rightarrow \infty} \Sigma(t)\psi = \mathcal{E}_0$ .
- (ii) If  $u_1^0 \not\equiv 0, u_2^0 \equiv 0$ , then we have  $u_1(\cdot, t, \psi) > 0$  and  $u_2(\cdot, t, \psi) \equiv 0$ . Then the equations for  $(R, S, u_1)$  satisfy system (3.4)–(3.6) with  $i = 1$ . Since  $\lambda_1^0 > 0$ , it follows from Lemma 3.2 that  $\lim_{t \rightarrow \infty} (R(\cdot, t, \psi), S(\cdot, t, \psi), u_1(\cdot, t, \psi)) = (R_1^*(x), S_1^*(x), u_1^*(x))$ , and hence,  $\lim_{t \rightarrow \infty} \Sigma(t)\psi = \mathcal{E}_1$ .
- (iii) If  $u_1^0 \equiv 0, u_2^0 \not\equiv 0$ , then we can use the fact  $\lambda_2^0 > 0$  and the same arguments as in (ii) to show that  $\lim_{t \rightarrow \infty} \Sigma(t)\psi = \mathcal{E}_2$ .
- (iv) If  $u_1^0 \not\equiv 0, u_2^0 \not\equiv 0$ , then we have  $u_1(\cdot, t, \psi) > 0$  and  $u_2(\cdot, t, \psi) > 0$ . Since (3.15) holds, it follows from Theorem 3.1 and Remark 3.1 that

$$(R(\cdot, t, \psi), S(\cdot, t, \psi), u_1(\cdot, t, \psi), u_2(\cdot, t, \psi))$$

will eventually enter the global attractor  $A_0 \subset \text{Int}(C([0, 1], \mathbb{R}_+^4))$ . Thus,  $\Sigma(t)\psi$  will eventually enter the global attractor  $M_3$ .

In the case where  $Z(\cdot, t_1, \psi) \not\equiv 0$ , for some  $t_1 \geq 0$ . Then Lemma 2.3 implies that  $Z(x, t, \psi) > 0, \forall x \in [0, 1], \forall t > t_1$ . It then follows that for each  $t > t_1$ , either  $u_1(\cdot, t, \psi) \equiv 0$  or  $u_2(\cdot, t, \psi) \equiv 0$ . If  $u_1(\cdot, t, \psi) \equiv 0$ , for each  $t > t_1$ . Then  $G(u_1(\cdot, t, \psi), u_2(\cdot, t, \psi)) = 0$ , and hence,  $Z(x, t, \psi)$  satisfies

$$\begin{cases} \frac{\partial Z}{\partial t} = d \frac{\partial^2 Z}{\partial x^2}, & x \in (0, 1), \quad t > 0, \\ \frac{\partial Z}{\partial x}(0, t) = \frac{\partial Z}{\partial x}(1, t) + \gamma Z(1, t) = 0, & t > 0, \\ Z(x, 0) = Z^0(x) \geq 0, & x \in (0, 1). \end{cases} \tag{4.18}$$

It is easy to see that  $\lim_{t \rightarrow \infty} Z(\cdot, t, \psi) = 0$ . Thus, either  $\lim_{t \rightarrow \infty} \Sigma(t)\psi = \mathcal{E}_0$  or  $\lim_{t \rightarrow \infty} \Sigma(t)\psi = \mathcal{E}_2$ . If  $u_1(\cdot, t_2, \psi) \not\equiv 0$ , for some  $t_2 > t_1$ . Then Lemma 2.3 implies that  $u_1(x, t, \psi) > 0, \forall x \in [0, 1], \forall t > t_2$ . It then follows that for each  $t > t_2, u_2(\cdot, t, \psi) \equiv 0$ . Thus,  $G(u_1(\cdot, t, \psi), u_2(\cdot, t, \psi)) = 0$ , and hence,  $Z(x, t, \psi)$  satisfies (4.18). This implies that either  $\lim_{t \rightarrow \infty} \Sigma(t)\psi = \mathcal{E}_0$  or  $\lim_{t \rightarrow \infty} \Sigma(t)\psi = \mathcal{E}_1$ . we have proved claim 1.

*Claim 2.* For  $i = 0, 1, 2, 3, M_i$  is a uniform weak repeller for  $\mathbb{X}_0$  in the sense that there exists  $\delta_i > 0$  such that  $\limsup_{t \rightarrow \infty} \|\Sigma(t)\mathbf{v}^0 - M_i\| \geq \delta_i, \forall \mathbf{v}^0 \in \mathbb{X}_0$ .

By Lemma 4.2, it follows that claim 2 is true when  $i = 3$ . Next, we only give the detailed arguments for the case  $i = 2$  since we can prove the cases where  $i = 0, 1$  by using the similar arguments. From the

fact that  $\Lambda_2^0 > 0$ , we may assume that there exists an  $\epsilon_2 > 0$  such that  $\Lambda_{2\epsilon}^0 > 0$ , where  $\Lambda_{2\epsilon}^0$  is the principal eigenvalue of the eigenvalue problem

$$\begin{cases} d\varphi''(x) + [f_1(R_2^*(x), S_2^*(x))e^{-\alpha u_2^*(x)} - 2\epsilon_2]\varphi(x) = \Lambda_2\varphi(x), & x \in (0, 1), \\ \varphi'(0) = \varphi'(1) + \gamma\varphi(1) = 0, \end{cases}$$

where  $R_2^*(x) := w_R(x) - q_{2r}u_2^*(x)$  and  $S_2^*(x) := w_S(x) - q_{2s}u_2^*(x)$  (see (3.9)). The positive eigenfunction corresponding to  $\Lambda_{2\epsilon}^0$  can be uniquely determined by the normalization, and we denote it by  $\varphi_{2\epsilon}^0(x)$ .

It is easy to see that there exists  $\delta_{21} > 0$  such that if  $\|(R_2, S_2, u_2) - (R_2^*(\cdot), S_2^*(\cdot), u_2^*(\cdot))\| < \delta_{21}$ , then

$$f_1(R_2, S_2)e^{-\alpha u_2} > f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - \epsilon_2. \tag{4.19}$$

Rewrite  $q_1g_1(u_1)Z = I(u_1, Z)u_1$ , where  $I(u_1, Z) = q_1 \frac{g_1(u_1)}{u_1}Z$  and  $I(0, 0) = 0$ . Then there exists  $\delta_{22} > 0$  such that if  $\|(u_1, Z) - (0, 0)\| < \delta_{22}$ , then

$$I(u_1, Z) < I(0, 0) + \epsilon_2 = \epsilon_2. \tag{4.20}$$

Setting  $\delta_2 := \min\{\delta_{21}, \delta_{22}\}$ . Then we show that

$$\limsup_{t \rightarrow \infty} \|\Sigma(t)\mathbf{v}^0 - M_2\| \geq \delta_2, \quad \forall \mathbf{v}^0 \in \mathbb{X}_0. \tag{4.21}$$

Suppose, by contradiction, there exists  $\mathbf{v}_0^0 \in \mathbb{X}_0$  such that  $\limsup_{t \rightarrow \infty} \|\Sigma(t)\mathbf{v}_0^0 - M_2\| < \delta_2$ . Then there exists  $\tilde{t} > 0$  such that for  $t \geq \tilde{t}$  and  $x \in [0, 1]$ , we have

$$\|(R(x, t, \mathbf{v}_0^0), S(x, t, \mathbf{v}_0^0), u_2(x, t, \mathbf{v}_0^0)) - (R_2^*(x), S_2^*(x), u_2^*(x))\| < \delta \leq \delta_{21},$$

and

$$\|(u_1(x, t, \mathbf{v}_0^0), Z(x, t, \mathbf{v}_0^0)) - (0, 0)\| < \delta \leq \delta_{22}.$$

It follows from (4.19) and (4.20) that for  $t \geq \tilde{t}$ , we have

$$f_1(R(\cdot, t, \mathbf{v}_0^0), S(\cdot, t, \mathbf{v}_0^0))e^{-\alpha u_2(\cdot, t, \mathbf{v}_0^0)} > f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - \epsilon_2, \tag{4.22}$$

and

$$I(u_1(\cdot, t, \mathbf{v}_0^0), Z(\cdot, t, \mathbf{v}_0^0)) < \epsilon_2. \tag{4.23}$$

With (4.22) and (4.23), it follows from the third equation of (1.1) that

$$\begin{cases} \frac{\partial u_1}{\partial t} \geq d \frac{\partial^2 u_1}{\partial x^2} + [f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - 2\epsilon_2]u_1, & x \in (0, 1), \quad t \geq \tilde{t}, \\ \frac{\partial u_1}{\partial x}(0, t) = \frac{\partial u_1}{\partial x}(1, t) + \gamma u_1(1, t) = 0, & t \geq \tilde{t}. \end{cases} \tag{4.24}$$

Since  $\mathbf{v}_0^0 \in \mathbb{X}_0$ , it follows from Lemma 2.3 that  $u_1(x, t, \mathbf{v}_0^0) > 0, \forall x \in [0, 1], t > 0$ . Thus, there exists a sufficiently small number  $\rho_0 > 0$  such that  $u_1(x, \tilde{t}, \mathbf{v}_0^0) \geq \rho_0 \varphi_{2\epsilon}^0(x), \forall x \in [0, 1]$ . Note that  $\tilde{u}_1(x, t) := \rho_0 e^{\Lambda_{2\epsilon}^0(t-\tilde{t})} \varphi_{2\epsilon}^0(x), t \geq \tilde{t}$ , is a solution of the following linear system:

$$\begin{cases} \frac{\partial u_1}{\partial t} = d \frac{\partial^2 u_1}{\partial x^2} + [f_1(R_2^*(\cdot), S_2^*(\cdot))e^{-\alpha u_2^*(\cdot)} - 2\epsilon_2]u_1, & x \in (0, 1), \quad t \geq \tilde{t}, \\ \frac{\partial u_1}{\partial x}(0, t) = \frac{\partial u_1}{\partial x}(1, t) + \gamma u_1(1, t) = 0, & t \geq \tilde{t}, \end{cases} \tag{4.25}$$



with initial data  $\tilde{u}_1(x, \tilde{t}) := \rho_0 \varphi_{2\epsilon}^0(x)$ . Then the comparison principle implies that

$$u_1(x, t, \mathbf{v}_0^0) \geq \tilde{u}_1(x, t) := \rho_0 e^{\Lambda_{2\epsilon}^0(t-\tilde{t})} \varphi_{2\epsilon}^0(x), \quad \forall x \in [0, 1], t \geq \tilde{t}.$$

Since  $\Lambda_{2\epsilon}^0 > 0$ , it follows that  $u_1(x, t, \mathbf{v}_0^0)$  is unbounded. This contradiction proves the result in (4.21). Thus, claim 2 holds.

Define a continuous function  $\mathbf{p} : \mathbb{X} \rightarrow [0, \infty)$  by

$$\mathbf{p}(\mathbf{v}^0) := \min_{3 \leq i \leq 5} \{ \min_{x \in [0,1]} v_i^0(x) \}, \quad \forall \mathbf{v}^0 := (v_1^0, v_2^0, v_3^0, v_4^0, v_5^0) \in \mathbb{X}.$$

By Lemma 2.3, it follows that  $\mathbf{p}^{-1}(0, \infty) \subseteq \mathbb{X}_0$  and  $\mathbf{p}$  has the property that if  $\mathbf{p}(\mathbf{v}^0) > 0$  or  $\mathbf{v}^0 \in \mathbb{X}_0$  with  $\mathbf{p}(\mathbf{v}^0) = 0$ , then  $\mathbf{p}(\Sigma(t)\mathbf{v}^0) > 0, \forall t > 0$ . That is,  $\mathbf{p}$  is a generalized distance function for the semiflow  $\Sigma(t) : \mathbb{X} \rightarrow \mathbb{X}$  (see, e.g., [15]). By the above claims, it follows that any forward orbit of  $\Sigma(t)$  in  $M_\partial$  converges to either  $M_0$  or  $M_1$  or  $M_2$  or  $M_3$ . Further,  $M_0, M_1, M_2,$  and  $M_3$  are isolated in  $\mathbb{X}$  and  $W^s(M_i) \cap \mathbb{X}_0 = \emptyset, \forall i = 0, 1, 2, 3$ , where  $W^s(M_i)$  is the stable set of  $M_i, i = 0, 1, 2, 3$  (see [15]). It is easy that no subsets of  $M_0, M_1, M_2, M_3$  forms a cycle in  $M_\partial$ .

By Lemma 2.2, it is easy to see that  $\Sigma(t) : \mathbb{X} \rightarrow \mathbb{X}$  has a global compact attractor in  $\mathbb{X}, \forall t \geq 0$ . It follows from [15, Theorem 3] that there exists an  $\eta > 0$  such that

$$\min_{\psi \in \omega(\mathbf{v}^0)} \mathbf{p}(\psi) > \eta, \quad \forall \mathbf{v}^0 \in \mathbb{X}_0.$$

This implies that (4.17) holds. Hence, the uniform persistence stated in our theorem is valid. By [9, Theorem 3.7 and Remark 3.10], it then follows that  $\Sigma(t) : \mathbb{X}_0 \rightarrow \mathbb{X}_0$  has a global attractor. It then follows from [9, Theorem 4.7] that  $\Sigma(t)$  has an steady-state solution  $(\hat{R}(\cdot), \hat{S}(\cdot), \hat{u}_1(\cdot), \hat{u}_2(\cdot), \hat{Z}(\cdot)) \in \mathbb{X}_0. \square$

### 5. Discussion

In this paper, we propose and analyze an unstirred chemostat model of the dynamics of *P. parvum*, cyanobacteria, and a zooplankton population. In our system (1.1)–(1.3), *P. parvum* competes for nutrients with cyanobacteria, which inhibits the growth of *P. parvum*. The zooplankton population grazes on *P. parvum* and cyanobacteria for growth, but *P. parvum* also inhibits the growth of zooplankton. This project is highly motivated by paper [6], in which the authors investigated a well-mixed chemostat system modeling the inhibitory/allelopathic effects of the algal toxins produced by *P. parvum* and cyanobacteria. Our system (1.1)–(1.3) further includes spatial variations, but neglects the compartments of algal toxins produced by *P. parvum* and cyanobacteria. The strength of inhibition/allelopathy is directly determined by the densities of *P. parvum* and cyanobacteria, respectively, not their toxins, which reduces the numbers of the modeling equations.

In order to study the coexistence of system (1.1)–(1.3), we need first to find the following possible steady-state solutions: the trivial steady-state solution of (1.1)–(1.3), corresponds to the absence of both harmful algae and zooplankton, is unique (Lemma 3.2 (i)); two semi-trivial steady-state solutions, corresponds to the presence of one of the algae and the absence of the other algae and zooplankton, are both unique if they exist (Lemma 3.2 (ii)); zooplankton-extinct steady-state solution, corresponds to the presence of both harmful algae and the absence of zooplankton, is not necessarily unique (see Theorem 3.1 and Remark 3.1). After defining a suitable continuous function  $m(x)$  (see Lemma 4.1), we are able to show that the compact attractor  $M_3$  on the boundary  $Z = 0$  is a uniform weak repeller for system (1.1)–(1.3) (see Lemma 4.2) under appropriate conditions. Then, we are able to show that system (1.1)–(1.3) is uniformly persistent, and system (1.1)–(1.3) admits at least one (componentwise) positive steady-state solution when the trivial

steady-state solution, two semi-trivial steady-state solutions, and the compact set  $M_3$  are all invisable (see Theorem 4.1).

An important problem is about the realization of the condition (3.15) and  $\mu^0 > 0$ . In view of (3.8), (3.11) and (3.13), it follows that  $\lambda_i^0$ ,  $\Lambda_1^0$ , and  $\Lambda_2^0$  can be characterized as (see, e.g., [1])

$$\lambda_i^0 = \sup_{\varphi \neq 0, \varphi \in H} \frac{\int_0^1 f_i(w_R(x), w_S(x)) \varphi^2 dx - d[\gamma(\varphi(1))^2 + \int_0^1 \varphi_x^2 dx]}{\int_0^1 \varphi^2 dx}, \quad i = 1, 2, \quad (5.1)$$

$$\Lambda_1^0 = \sup_{\varphi \neq 0, \varphi \in H} \frac{\int_0^1 f_2(R_1^*(x), S_1^*(x)) \varphi^2 dx - d[\gamma(\varphi(1))^2 + \int_0^1 \varphi_x^2 dx]}{\int_0^1 \varphi^2 dx}, \quad (5.2)$$

and

$$\Lambda_2^0 = \sup_{\varphi \neq 0, \varphi \in H} \frac{\int_0^1 f_1(R_2^*(x), S_2^*(x)) e^{-\alpha u_2^*(x)} \varphi^2 dx - d[\gamma(\varphi(1))^2 + \int_0^1 \varphi_x^2 dx]}{\int_0^1 \varphi^2 dx}, \quad (5.3)$$

where  $H$  represents the closure of  $C^1([0, 1])$  with respect to the topology induced by the norm

$$\|\theta\| = \left( \int_0^1 \theta^2 dx + \int_0^1 \theta_x^2 dx \right)^{1/2}.$$

With (5.1), (5.2) and (5.3), we may expect that (3.15) holds if the diffusion rate  $d > 0$  is small. Similarly, it follows from (1.4), (1.5), (4.3) and (4.7) that we can also expect that  $\mu^0 > 0$  if both  $m_{1z}$  and  $m_{2z}$  are sufficiently large. However, it remains a challenging task to find an exact parameter region in which (3.15) and  $\mu^0 > 0$  are valid.

Our work extended the well-mixed model in [6] to a partially-mixed system, but we also did two significant simplifications in modeling. For example, the growth rate of zooplankton  $G(u_1, u_2)$  only takes the form (1.4) or (1.5), which eliminates the possibility of two types of steady-state solutions for system (1.1)–(1.3) (see (4.1) and (4.2)). If  $G(u_1, u_2)$  takes the substitutable type, that is,  $G(0, u_2)$  and  $G(u_1, 0)$  can be both positive, then it is likely that another two types of steady-state solutions, (4.1) and (4.2), can happen. This will make the analysis much more complicated than those in the current paper. On the other hand, in order to reduce the numbers of model equations in our system, we have removed the compartments of algal toxins produced by both harmful algae, and the associated inhibition/allelopathy is directly affected by the densities of algae respectively. We will relax the aforementioned simplifications and investigate a more realistic and challenging case in the future.

## Acknowledgment

We are grateful to the anonymous referee for careful reading and helpful suggestions which led to improvements of our original manuscript.

## References

- [1] R.S. Cantrell, C. Cosner, *Spatial Ecology via Reaction–Diffusion Equations*, Wiley Ser. Math. Comput. Biol., John Wiley and Sons, Chichester, UK, 2003.
- [2] J. Hale, *Asymptotic Behavior of Dissipative Systems*, American Mathematical Society, Providence, RI, 1988.
- [3] S.B. Hsu, P. Waltman, On a system of reaction–diffusion equations arising from competition in an unstirred chemostat, *SIAM J. Appl. Math.* 53 (1993) 1026–1044.
- [4] F.X. Fu, A.O. Tatters, D.A. Hutchins, Global change and the future of harmful algal blooms in the ocean, *Mar. Ecol. Prog. Ser.* 470 (2012) 207–233.

- [5] J.P. Grover, K.W. Crane, J.W. Baker, B.W. Brooks, D.L. Roelke, Spatial variation of harmful algae and their toxins in flowing-water habitats: a theoretical exploration, *J. Plankton Res.* 33 (2011) 211–227.
- [6] J.P. Grover, D.L. Roelke, B.W. Brooks, Modeling of plankton community dynamics characterized by algal toxicity and allelopathy: a focus on historical *Prymnesium parvum* blooms in a Texas reservoir, *Ecol. Model.* 227 (2012) 147–161.
- [7] G.M. Hallegraeff, A review of harmful algal blooms and their apparent global increase, *Phycologia* 32 (1993) 79–99.
- [8] R. Martin, H.L. Smith, Abstract functional differential equations and reaction–diffusion systems, *Trans. Amer. Math. Soc.* 321 (1990) 1–44.
- [9] P. Magal, X.-Q. Zhao, Global attractors and steady states for uniformly persistent dynamical systems, *SIAM J. Math. Anal.* 37 (2005) 251–275.
- [10] H. Nie, J. Wu, G.S.K. Wolkowicz, A mathematical model of competition for two essential resources in the unstirred chemostat, *SIAM J. Appl. Math.* 65 (2004) 209–229.
- [11] A. Pazy, *Semigroups of Linear Operators and Application to Partial Differential Equations*, Springer-Verlag, 1983.
- [12] M.H. Protter, H.F. Weinberger, *Maximum Principles in Differential Equations*, Springer-Verlag, 1984.
- [13] H.L. Smith, *Monotone Dynamical Systems: An Introduction to the Theory of Competitive and Cooperative Systems*, *Math. Surveys Monogr.*, vol. 41, American Mathematical Society, Providence, RI, 1995.
- [14] H.L. Smith, P.E. Waltman, *The Theory of the Chemostat*, Cambridge Univ. Press, 1995.
- [15] H.L. Smith, X.-Q. Zhao, Robust persistence for semidynamical systems, *Nonlinear Anal.* 47 (2001) 6169–6179.
- [16] F.B. Wang, A system of partial differential equations modeling the competition for two complementary resources in flowing habitats, *J. Differential Equations* 249 (2010) 2866–2888.
- [17] F.-B. Wang, C.-C. Huang, A reaction–advection–diffusion system modeling the competition for two complementary resources with seasonality in a flowing habitat, *J. Math. Anal. Appl.* 428 (2015) 145–164.
- [18] J. Wu, G.S.K. Wolkowicz, A system of resource-based growth models with two resources in the unstirred chemostat, *J. Differential Equations* 172 (2001) 300–332.
- [19] X.-Q. Zhao, *Dynamical Systems in Population Biology*, Springer, New York, 2003.