

## SET 6

### MATH 543: FROBENIUS METHOD

References: Hildebrand and Sadri Hassan.

The following theorem summarizes the Frobenius method: (Proved in Class please see your lecture notes and also DK)

**Theorem 1.** *Suppose that the differential equation  $Lu = 0$ , where  $L$  is a second order differential operator, has a regular singular point at  $z = z_0$  with the roots  $r_1$  and  $r_2$  of the indicial equation. There are three possible cases: (Assuming  $\operatorname{Re}(r_1) > \operatorname{Re}(r_2)$ )*

1.  $r_1 - r_2 \neq \text{an integer}$
2.  $r_1 - r_2 = N$  (a non-negative integer) and recursion relation is consistent
3.  $r_1 - r_2 = N$  (a non-negative integer) and recursion relation is not consistent.

*Then, in the first two cases, there exists a bases of  $\{u_1, u_2\}$  of solutions of  $Lu = 0$ . These solutions are of the form*

$$u_1(z) = (z - z_0)^{r_1} \sum_{k=0}^{\infty} C_k^1 (z - z_0)^k, \quad (1)$$

$$u_2(z) = (z - z_0)^{r_2} \sum_{k=0}^{\infty} C_k^2 (z - z_0)^k \quad (2)$$

*and the third case, the bases  $\{u_1, u_2\}$  is of the following form*

$$u_1(z) = (z - z_0)^{r_1} \sum_{k=0}^{\infty} C_k (z - z_0)^k, \quad (3)$$

$$u_2(z) = C u_1(z) \ln(z - z_0) + (z - z_0)^{r_2} \sum_{k=1}^{\infty} B_k (z - z_0)^k \quad (4)$$

*where the power series about  $(z - z_0)$  are convergent in a neighborhood of  $z = z_0$*

**A method when  $r_1 - r_2 = N$**

Let  $z = z_0$  be a regular singular point of differential equation

$$Lu = u'' + p(z)u' + q(z)u = 0, \quad (5)$$

where

$$A(z) = (z - z_0)p(z), \quad B(z) = (z - z_0)^2 q(z) \quad (6)$$

are analytic in a neighborhood  $D$  of  $z = z_0$  hence  $A(z)$  and  $B(z)$  are analytic in  $D$  then

$$A(z) = \sum_{k=0}^{\infty} a_k (z - z_0)^k, \quad B(z) = \sum_{k=0}^{\infty} b_k (z - z_0)^k, \quad (7)$$

Using the anzats

$$u(z) = (z - z_0)^r \sum_{k=0}^{\infty} C_k (z - z_0)^k, \quad (8)$$

in the DE we get

$$\lambda_0(n + r) C_n + \sum_{m=0}^{n-1} [(m + r)a_{n-m} + b_{n-m}] C_m, \quad (9)$$

This is the recursion relations mentioned in the above theorem 1. Here

$$\lambda_0(r) = r(r - 1) + r a_0 + b_0 \quad (10)$$

If the indicial equation is not imposed  $\lambda_0(r) \neq 0$  and the remaining

$$C_1(r), C_2(r), \dots, C_n(r) \quad (11)$$

are determined through the recursion relation. Since  $n = 0$  term is not set to zero then the differential equation is not satisfied but becomes

$$Lu = C_0 \lambda_0(r) (z - z_0)^{r-2} \quad (12)$$

where  $C_0$  is the undetermined constant. The solution of this equation is

$$u(r, z) = (z - z_0)^r \sum_{k=0}^{\infty} C_k(r) (z - z_0)^k, \quad (13)$$

In the case 3 , when the difference of the indices is a nonegative number and recursion relations are inconsistent the the usual anzats fails. To find the correct anzats and a method of solution we use (12) and (13).

$r_1 = r_2$  **case**

Eq.(12) becomes

$$Lu = C_0 (r - r_1)^2 (z - z_0)^{r-2}, \quad (14)$$

Taking the derivative of both sides wrt  $r$  and letting  $r = r_1$  we obtain the second solution as

$$u_2(z) = \frac{du(r, z)}{dr} \Big|_{r=r_1}, \quad (15)$$

Using the form of  $u(r, z)$  given in (13) we obtain

$$u_2(z) = u_1(z) \ln(z - z_0) + (z - z_0)^{r_1} \sum_{k=0}^{\infty} C'_k(r_1) (z - z_0)^k \quad (16)$$

where  $C'_k(r_1) = \frac{d}{dr} C_k(r) \Big|_{r=r_1}$

$r_1 - r_2 = N$  **a positive integer case**

Eq.(12) becomes

$$Lu = C_0(r - r_1)(r - r_2) (z - z_0)^{r-2} \quad (17)$$

Multiplying both sides by  $r - r_2$  and taking derivatives of both sides wrt  $r$  and taking limit as  $r$  goes to  $r_2$  we obtain

$$u_2(z) = \frac{d}{dr} [(r - r_2) u(r, z)] \text{ at } r = r_2 \quad (18)$$

Using the expression (13) for  $u(r, z)$  we obtain

$$u_2(z) = Cu_1(z) \ln(z - z_0) + (z - z_0)^{r_2} \sum_{k=0}^{\infty} \tilde{C}_k (z - z_0)^k \quad (19)$$

where

$$\tilde{C}_k = \frac{d}{dr}[(r - r_2) C_k(r)]|_{r=r_2}$$

### Problems

1. An example for the case  $r_1 - r_2 = N$  with consistent recursion relations. Solve  $z^2 u'' + (z^2 + z)u' - u = 0$  about  $z = 0$  (solved in Class)
2. An example for the case  $r_1 = r_2$  with inconsistent recursion relations. Solve  $z^2 u'' + zu' + z^2 u = 0$  about  $z = 0$ . You can find the solution  $u_1$  by using the recursion relations. This solution is known as the Bessel function of order 0,

$$u_1(z) = J_0(z) = \sum_{k=0}^{\infty} \frac{(-1)^k (z/2)^{2k}}{(k!)^2}$$

The second solution  $u_2(z)$  is found by using (16). Prove that it takes the form

$$u_2(z) = J_0(z) \ln z + \sum_{k=0}^{\infty} \varphi(k) \frac{(z/2)^{2k}}{(k!)^2}$$

where

$$\varphi(k) = \begin{cases} \sum_{m=1}^k \frac{1}{m}, & \text{for } k = 1, 2, \dots \\ 0, & \text{for } (k = 0) \end{cases} \quad (20)$$

3. An example for the case  $r_1 - r_2 = N$  with inconsistent recursion relations. Solve  $z u'' - u = 0$  about  $z = 0$ . Again  $u_1(z)$  will be found by the standard method

$$u_1(z) = \sum_{k=0}^{\infty} \frac{z^{k+1}}{k!(k+1)!}$$

And the second solution is found by using the formula (19)

$$u_2(z) = u_1 \ln z + 1 - \sum_{k=1}^{\infty} \frac{\varphi(k) + \varphi(k-1)}{k!(k-1)!} z^k$$

4. Find the solutions of the Bessel's equation  $z^2 u'' + zu' + (z^2 - p^2)u = 0$ . Solutions of this equation are known as the Bessel's functions of order  $p$ . Solve this equation for  $p = \text{not integer}$  and for  $p = \text{integer}$ .

5. Show that for the differential equation  $zu'' + 3u' + 4zu = 0$  the condition  $u(0) = 1$  determines a unique solution, and that  $u'(0)$  cannot also be prescribed. Determine this solution.

6. Use the method of Frobenius to obtain the general solution of each of the following differential equations, valid near  $z = 0$ :

(1)  $z^2u'' - 2zu' + (2 - z^2)u = 0$

(2)  $(z - 1)u'' - zu' + u = 0$

(3)  $2zu'' + (1 - 2z)u' - u = 0$

(4)  $z^2u'' + zu' + (z^2 - \frac{1}{4})u = 0$

(5)  $zu'' - u' + 4z^3u = 0$

(6)  $zu'' + 2u' + zu = 0$

(7)  $z(1 - z)u'' - 2u' - 2 + 2u = 0$

7. Determine the two values of the constant  $\alpha$  for which it is true that all solutions of the equation  $zu'' + (z - 1)u' - \alpha u = 0$  are regular at  $z = 0$ , and obtain the general solution in each of these cases

8. (a) Show that the equation  $zu'' + u' - u = 0$  possesses equal indices  $r_1 = r_2 = 0$  at  $z = 0$

(b) Obtain the regular solution  $u_1(z)$

(c) Assume the second solution of the form  $u_2(z) = Cu_1(z) \ln z + v(z)$  where  $C \neq 0$ . Find  $v(z)$ .

9. (a) Show that the equation  $zu'' - zu' - u = 0$  possesses indices  $r_1 = 1, r_2 = 0$  at  $z = 0$ .

(b) Obtain the regular solution  $u_1(z)$

(c) Assume the general form for the second solution as  $u_2(z) = Cu_1 \ln z + v(z)$ , where  $C \neq 0$ . Find  $v(z)$ .