

1.
  - (a) Intersecting lines with solution  $x_1 = x_2 = 1$ .
  - (b) One line, so there is an infinite number of solutions with  $x_2 = \frac{3}{2} - \frac{1}{2}x_1$ .
  - (c) One line, so there is an infinite number of solutions with  $x_2 = -\frac{1}{2}x_1$ .
  - (d) Intersecting lines with solution  $x_1 = \frac{2}{7}$  and  $x_2 = -\frac{11}{7}$ .
2.
  - (a) Intersecting lines whose solution is  $x_1 = x_2 = 0$ .
  - (b) Parallel lines, so there is no solution.
  - (c) Three lines in the plane that do not intersect at a common point.
  - (d) Two planes in space which intersect in a line with  $x_1 = -\frac{5}{4}x_2$  and  $x_3 = \frac{3}{2}x_2 + 1$ .
3. Gaussian elimination gives the following solutions.
  - (a)  $x_1 = 1.0, x_2 = -0.98, x_3 = 2.9$
  - (b)  $x_1 = 1.1, x_2 = -1.1, x_3 = 2.9$
4. Gaussian elimination gives the following solutions.
  - (a)  $x_1 = -0.70, x_2 = 1.1, x_3 = 2.9$
  - (b)  $x_1 = -0.88, x_2 = 0.74, x_3 = 3.0$
5. Gaussian elimination gives the following solutions.
  - (a)  $x_1 = 1.1875, x_2 = 1.8125, x_3 = 0.875$  with one row interchange required
  - (b)  $x_1 = -1, x_2 = 0, x_3 = 1$  with no interchange required
  - (c)  $x_1 = 1.5, x_2 = 2, x_3 = -1.2, x_4 = 3$  with no interchange required
  - (d) No unique solution
6. Gaussian elimination gives the following solutions.
  - (a)  $x_1 = -4, x_2 = -8, x_3 = -6$  with one row interchange required
  - (b)  $x_1 = \frac{22}{9}, x_2 = -\frac{4}{9}, x_3 = \frac{4}{3}, x_4 = 1$  with one row interchange required
  - (c)  $x_1 = 13, x_2 = 8, x_3 = 8, x_4 = 5$  with one row interchange required.
  - (d)  $x_1 = -1, x_2 = 2, x_3 = 0, x_4 = 1$  with one row interchange required.
7. Gaussian elimination with *Digits:=10* gives the following solutions:
  - (a)  $x_1 = -227.0769, x_2 = 476.9231, x_3 = -177.6923$ ;
  - (b)  $x_1 = 1.001291, x_2 = 1, x_3 = 1.00155$ ;
  - (c)  $x_1 = -0.03174600, x_2 = 0.5952377, x_3 = -2.380951, x_4 = 2.777777$ ;
  - (d)  $x_1 = 1.918129, x_2 = 1.964912, x_3 = -0.9883041, x_4 = -3.192982, x_5 = -1.134503$ .

8. Gaussian elimination with  $Digits:=10$  gives the following solutions:

- (a)  $x_1 = 0.9798657720, x_2 = 4.281879191, x_3 = 17.48322147;$
- (b)  $x_1 = 6.461447620, x_2 = 8.394321092, x_3 = -0.01347368618;$
- (c)  $x_1 = 1.349448559, x_2 = -4.67798776, x_3 = -4.032893779, x_4 = -1.656637732;$
- (d)  $x_1 = 13.49999998, x_2 = -11.5000000000, x_3 = 23.75000003, x_4 = 121.5000003, x_5 = 97.75000025.$

9. (a) When  $\alpha = -1/3$ , there is no solution.
- (b) When  $\alpha = 1/3$ , there is an infinite number of solutions with  $x_1 = x_2 + 1.5$ , and  $x_2$  is arbitrary.
- (c) If  $\alpha \neq \pm 1/3$ , then the unique solution is

$$x_1 = \frac{3}{2(1+3\alpha)} \quad \text{and} \quad x_2 = \frac{-3}{2(1+3\alpha)}.$$

10. (a)  $\alpha = 1$
- (b)  $\alpha = -1$
- (c)  $x_1 = -1/(1-\alpha), x_2 = 1, x_3 = 1/(1-\alpha)$

11. Suppose  $x'_1, \dots, x'_n$  is a solution to the linear system (6.1).

(i) The new system becomes

$$\begin{aligned} E_1 : a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\ &\vdots \\ E_i : \lambda a_{i1}x_1 + \lambda a_{i2}x_2 + \dots + \lambda a_{in}x_n &= \lambda b_i \\ &\vdots \\ E_n : a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n &= b_n. \end{aligned}$$

Clearly,  $x'_1, \dots, x'_n$  satisfies this system. Conversely, if  $x_1^*, \dots, x_n^*$  satisfies the new system, dividing  $E_i$  by  $\lambda$  shows  $x_1^*, \dots, x_n^*$  also satisfies (6.1).

(ii) The new system becomes

$$\begin{aligned} E_1 : a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\ &\vdots \\ E_i : (a_{i1} + \lambda a_{j1})x_1 + (a_{i2} + \lambda a_{j2})x_2 + \dots + (a_{in} + \lambda a_{jn})x_n &= b_i + \lambda b_j \\ &\vdots \\ E_n : a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n &= b_n. \end{aligned}$$

Clearly,  $x'_1, \dots, x'_n$  satisfies all but possibly the  $i$ th equation. Multiplying  $E_j$  by  $\lambda$  gives

$$\lambda a_{j1}x'_1 + \lambda a_{j2}x'_2 + \dots + \lambda a_{jn}x'_n = \lambda b_j,$$

which can be subtracted from  $E_i$  in the new system results in the system (6.1). Thus,  $x'_1, \dots, x'_n$  satisfies the new system. Conversely, if  $x_1^*, \dots, x_n^*$  is a solution to the new system, then all but possibly  $E_i$  of (6.1) are satisfied by  $x_1^*, \dots, x_n^*$ . Multiplying  $E_j$  of the new system by  $-\lambda$  gives

$$-\lambda a_{j1}x_1^* - \lambda a_{j2}x_2^* - \dots - \lambda a_{jn}x_n^* = -\lambda b_j.$$

Adding this to  $E_i$  in the new system produces  $E_i$  of (6.1). Thus,  $x_1^*, \dots, x_n^*$  is a solution of (6.1).

(iii) The new system and the old system have the same set of equations to satisfy. Thus, they have the same solution set.

12. Change Algorithm 6.1 as follows:

*STEP 1* For  $i = 1, \dots, n$  do STEPS 2, 3, and 4.

*STEP 4* For  $j = 1, \dots, i-1, i+1, \dots, n$  do STEPS 5 and 6.

*STEP 8* For  $i = 1, \dots, n$  set  $x_i = a_{i,n+1}/a_{ii}$ .

In addition, delete STEP 9.

13. The Gauss-Jordan method gives the following results.

(a)  $x_1 = 0.98, x_2 = -0.98, x_3 = 2.9$

(b)  $x_1 = 1.1, x_2 = -1.0, x_3 = 2.9$

14. The Gauss-Jordan method with single precision arithmetic gives the following solutions.

(a)  $x_1 = -227.0787, x_2 = 476.9262, x_3 = -177.6934$

(b)  $x_1 = 1.000036, x_2 = 0.9999991, x_3 = 0.9986052$

(c)  $x_1 = -0.03177120, x_2 = 0.5955572, x_3 = -2.381768, x_4 = 2.778329$

(d)  $x_1 = 1.918129, x_2 = 1.964912, x_3 = -0.9883036, x_4 = -3.192982, x_5 = -1.134503$

15. The results for are listed in the following table. (The abbreviations M/D and A/S are used for multiplications/divisions and additions/subtractions, respectively.)

| $n$ | Gaussian elimination |        | Gauss-Jordan |        |
|-----|----------------------|--------|--------------|--------|
|     | M/D                  | A/S    | M/D          | A/S    |
| 3   | 17                   | 11     | 21           | 12     |
| 10  | 430                  | 375    | 595          | 495    |
| 50  | 44150                | 42875  | 64975        | 62475  |
| 100 | 343300               | 338250 | 509950       | 499950 |

16. (a) The Gaussian elimination procedure requires

$$\frac{(2n^3 + 3n^2 - 5n)}{6} \text{ Multiplications/Divisions}$$

and

$$\frac{n^3 - n}{3} \text{ Additions/Subtractions.}$$

The additional elimination steps are:

$$\text{For } i = n, n-1, \dots, 2$$

$$\text{for } j = 1, \dots, i-1,$$

$$\text{set } a_{j,n+1} = a_{j,n+1} - \frac{a_{ji}a_{i,n+1}}{a_{ii}}.$$

This requires

$$n(n-1) \text{ Multiplications/Divisions}$$

and

$$\frac{n(n-1)}{2} \text{ Additions/Subtractions.}$$

Solving for

$$x_i = \frac{a_{i,n+1}}{a_{ii}}$$

requires  $n$  divisions. Thus, the totals are

$$\frac{n^3}{3} + \frac{3n^2}{2} - \frac{5n}{6} \text{ Multiplications/Divisions}$$

and

$$\frac{n^3}{3} + \frac{n^2}{2} - \frac{5n}{6} \text{ Additions/Subtractions.}$$

- (b) The results are listed in the following table. In this table the abbreviations M/D and A/S are used for Multiplications/Divisions and for Additions/Subtractions, respectively.

| $n$ | Gaussian Elimination |        | Gauss-Jordan |        | Hybrid |        |
|-----|----------------------|--------|--------------|--------|--------|--------|
|     | M/D                  | A/S    | M/D          | A/S    | M/D    | A/S    |
| 3   | 17                   | 11     | 21           | 12     | 20     | 11     |
| 10  | 430                  | 375    | 595          | 495    | 475    | 375    |
| 50  | 44150                | 42875  | 64975        | 62475  | 45375  | 42875  |
| 100 | 343300               | 338250 | 509950       | 499950 | 348250 | 338250 |

17. The Gaussian-Elimination-Gauss-Jordan hybrid method gives the following results.
- (a)  $x_1 = 1.0, x_2 = -0.98, x_3 = 2.9$
  - (b)  $x_1 = 1.0, x_2 = -1.0, x_3 = 2.9$
18. The Gauss-Jordan hybrid method with single-precision arithmetic gives the following solutions.
- (a)  $-227.0788, 476.9262, -177.6934$
  - (b)  $0.9990999, 0.9999991, 0.9986052$
  - (c)  $-0.03177060, 0.5955554, -2.381768, 2.778329$
  - (d)  $x_1 = 1.918126, x_2 = 1.964916, x_3 = -0.9883027, x_4 = -3.192982, x_5 = -1.134503$
19. (a) There is sufficient food to satisfy the average daily consumption.
- (b) We could add 200 of species 1, or 150 of species 2, or 100 of species 3, or 100 of species 4.
  - (c) Assuming none of the increases indicated in part (b) was selected, species 2 could be increased by 650, or species 3 could be increased by 150, or species 4 could be increased by 150.
  - (d) Assuming none of the increases indicated in parts (b) or (c) were selected, species 3 could be increased by 150, or species 4 could be increased by 150.

20. (a) For the Trapezoidal rule  $m = n = 1$ ,  $x_0 = 0$ ,  $x_1 = 1$  so that for  $i = 0$  and  $1$ , we have

$$\begin{aligned} u(x_i) &= f(x_i) + \int_0^1 K(x_i, t)u(t) dt \\ &= f(x_i) + \frac{1}{2} [K(x_i, 0)u(0) + K(x_i, 1)u(1)]. \end{aligned}$$

Substituting for  $x_i$  gives the desired equations.

- (b) We have  $n = 4$ ,  $h = \frac{1}{4}$ ,  $x_0 = 0$ ,  $x_1 = \frac{1}{4}$ ,  $x_2 = \frac{1}{2}$ ,  $x_3 = \frac{3}{4}$ , and  $x_4 = 1$ , so

$$\begin{aligned} u(x_i) &= f(x_i) + \frac{h}{2} \left[ K(x_i, 0)u(0) + 2K\left(x_i, \frac{1}{4}\right)u\left(\frac{1}{4}\right) \right. \\ &\quad \left. + 2K\left(x_i, \frac{1}{2}\right)u\left(\frac{1}{2}\right) + 2K\left(x_i, \frac{3}{4}\right)u\left(\frac{3}{4}\right) + K(x_i, 1)u(1) \right], \end{aligned}$$

for  $i = 0, 1, 2, 3, 4$ . This gives

$$u(x_i) = x_i^2 + \frac{1}{8} \left[ e^{x_i}u(0) + 2e^{|x_i - \frac{1}{4}|}u\left(\frac{1}{4}\right) + 2e^{|x_i - \frac{1}{2}|}u\left(\frac{1}{2}\right) + 2e^{|x_i - \frac{3}{4}|}u\left(\frac{3}{4}\right) + e^{|x_i - 1|}u(1) \right],$$

for each  $i = 1, \dots, 4$ . The  $5 \times 5$  linear system has solution  $u(0) = -1.154255$ ,  $u\left(\frac{1}{4}\right) = -0.9093298$ ,  $u\left(\frac{1}{2}\right) = -0.7153145$ ,  $u\left(\frac{3}{4}\right) = -0.5472949$ , and  $u(1) = -0.3931261$ .

- (c) The Composite Simpson's rule gives

$$\begin{aligned} \int_0^1 K(x_i, t)u(t) dt &= \frac{h}{3} \left[ K(x_i, 0)u(0) + 4K\left(x_i, \frac{1}{4}\right)u\left(\frac{1}{4}\right) + 2K\left(x_i, \frac{1}{2}\right)u\left(\frac{1}{2}\right) \right. \\ &\quad \left. + 4K\left(x_i, \frac{3}{4}\right)u\left(\frac{3}{4}\right) + K(x_i, 1)u(1) \right], \end{aligned}$$

which results in the linear equations

$$u(x_i) = x_i^2 + \frac{1}{12} \left[ e^{x_i}u(0) + 4e^{|x_i - \frac{1}{4}|}u\left(\frac{1}{4}\right) + 2e^{|x_i - \frac{1}{2}|}u\left(\frac{1}{2}\right) + 4e^{|x_i - \frac{3}{4}|}u\left(\frac{3}{4}\right) + e^{|x_i - 1|}u(1) \right].$$

The  $5 \times 5$  linear system has solutions  $u(0) = -1.234286$ ,  $u\left(\frac{1}{4}\right) = -0.9507292$ ,  $u\left(\frac{1}{2}\right) = -0.7659400$ ,  $u\left(\frac{3}{4}\right) = -0.5844737$ , and  $u(1) = -0.4484975$ .