

Midterm 01

Oct 24, 2017.

- (10 pts) The half precision format uses 16 bits to store a binary floating point number of the form $\pm 1.a_1a_2\cdots a_t \times 2^e$ where $a_j \in \{0, 1\}$, $-14 \leq e \leq 15$. Find t and derive an upper bound for relative error caused by rounding. Express your final answer as a real number, but need not convert it to decimal expression.
- (10 pts) Given p_0 , p_1 and p_2 , the general solution to the recursion formula $p_n = \frac{10}{3}p_{n-1} - 3p_{n-2} + \frac{2}{3}p_{n-3}$ is $p_n = c_11^n + c_22^n + c_3(\frac{1}{3})^3$ (need not show this). Find all $(c_1, c_2, c_3) \neq (0, 0, 0)$ such that the above iteration is unstable in relative error. Explain.
- (10 pts) The first few iteration $(p_i, f(p_i))$, $i = 0, 1, 2, 3, 4$ of method of false position for some equation $f(x) = 0$ is given by

$$(0, 2), \quad (3, 1), \quad (2, 2), \quad (1, 1), \quad \left(\frac{2}{3}, \frac{2}{9}\right)$$

Find p_5 (4 digits will do). Explain.

- (15 pts) Use any method to find a solution of $\sqrt{1+0.9x} - \sqrt{1-0.8x} = 1.0 \times 10^{-10}$ to 15 correct digits. You need to prevent loss of accuracy. Standard methods only gives you about 5 correct digits (and 1/3 partial credits).
- (10+5 pts) It is known that the unique solution to $f(x) = x + 3\sin(x) - 0.01 = 0$ is located near $x = 0$.
 - Find a fixed point iteration that will converge for any $x_0 \in [-\frac{1}{2}, \frac{1}{2}]$. Show that your method satisfies the assumptions of a relevant Theorem, but need not prove the Theorem again. You can use the numerical values of $\sin(\frac{1}{2})$, $\cos(\frac{1}{2})$, $\exp(\frac{1}{2})$, etc. in your proof.
 - Find an N (need not be optimal) such that $|x_n - x^*| < 10^{-30}$ for all $n \geq N$ with $x_0 = 0$ (assuming a higher precision floating point arithmetic is used).
- (15 pts) Give a cubically convergent method to solve for $e^x - 1 = 0$. Give the formula and prove that it is cubically convergent (locally). If you cannot do it, do the same for a locally quadratically convergent method for partial credit.
- (10 pts) Derive Aitken's Δ^2 acceleration method.
Hint: the starting point is to assume p_n converges to p linearly.
- (15 pts) Use any method to solve the nonlinear system of equations

$$\sin(x) + \frac{2y}{1+x} = 0.01, \quad 5x + \sin\left(\frac{6y}{1+y^2}\right) = 0.02.$$

Write your answer in the format of 'format long e'.

Hint: the solution is near $(0, 0)$ where $\sin x \approx x$, $\frac{y}{1+x} \approx y$, etc. to leading orders.