

Brief answers to Quiz 1

Oct 02, 2014

1. Give formal definition of $\lim_{x \rightarrow c} f(x) \neq L$ (just the definition, need not find ϵ or δ , etc.).

Ans:

See Remark 2 in ‘Study Guide for Chap 02 (New!’.

2. Find $\lim_{\theta \rightarrow 0} \frac{\sin(1 - \cos \theta)}{\tan^2 \theta}$.

Ans: 1/2. (See the hint in problem 3 of Homework 02)

3. State the Intermediate Value Theorem (Need not prove). Use it to show that $x^x = 2$ has a root.

Ans:

For the second part, apply Intermediate Value Theorem to the function $g(x) = x^x - 2$ on the interval $[1, 2]$.

4. Give formal definition of ‘ $y = f(x)$ is continuous at $x = c$ ’ in terms of ϵ and δ . Then use the $\epsilon - \delta$ argument to show that if both $f(x)$ and $g(x)$ are continuous at $x = c$, then so is $f(x) + g(x)$.

Ans:

The first part:

For any $\epsilon > 0$, there exists $\delta > 0$ such that,

$$|x - c| < \delta \implies |f(x) - f(c)| < \epsilon$$

The second part:

The proof is similar (and almost identical) to Example 6 of section 2.3. Just replace L and M by $f(c)$ and $g(c)$, respectively. See page 81 for details.

5. Give formal definitions of the following limits (Just the definition, need not find δ).

$$(a) \lim_{x \rightarrow c^-} f(x) = L \qquad (b) \lim_{x \rightarrow -\infty} f(x) = \infty$$

Ans:

- (a) For any $\epsilon > 0$, there exists $\delta > 0$ such that,

$$c - \delta < x < c \implies |f(x) - L| < \epsilon$$

- (b) For any $B > 0$, there exists $N < 0$ such that

$$x < N \implies f(x) > B$$