

Differentiation Rules

$$\text{Eg. } \frac{d}{dx} 1 = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$(f(x) = 1) = \lim_{h \rightarrow 0} \frac{1 - 1}{h} = 0$$

$$\text{Ex. } f(x) = x^2, \quad f'(x) = ?$$

$$\text{Ans. } \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = 2x$$

$$\text{Eg. } f(x) = \frac{1}{\sqrt{x}} = x^{-\frac{1}{2}}, \quad x > 0$$

$$f'(x) = \lim_{z \rightarrow x} \frac{\frac{1}{\sqrt{z}} - \frac{1}{\sqrt{x}}}{z - x} \cdot \frac{\frac{1}{\sqrt{z}} + \frac{1}{\sqrt{x}}}{\frac{1}{\sqrt{z}} + \frac{1}{\sqrt{x}}} = -\frac{1}{2} x^{-\frac{3}{2}}$$

Ex: $f(x) = x^n$. $n \in \mathbb{N}$
(positive integer)

$$\begin{aligned} f'(x) &= \lim_{z \rightarrow x} \frac{z^n - x^n}{z - x} \\ &= \lim_{z \rightarrow x} z^{n-1} + z^{n-2}x + \dots + x^{n-1} = nx^{n-1} \end{aligned}$$

These are examples of

$$\frac{d}{dx} x^n = nx^{n-1} \quad \dots \dots (*)$$

for $n = 0, 2, \frac{1}{2}, n \in \mathbb{N}$

Similarly, one can show that
(*) holds for $n \in \mathbb{Q}$

ie $n = \frac{q}{p}$, $p, q \in \mathbb{Z}$, $p \neq 0$

Example: $f(x) = x^{\frac{4}{3}}$,

$$f'(x) = \lim_{z \rightarrow x} \frac{z^{\frac{4}{3}} - x^{\frac{4}{3}}}{z - x}$$

$$((a-b)(a^2+ab+b^2) = a^3 - b^3)$$

$$= \lim_{z \rightarrow x} \frac{z^{\frac{4}{3}} - x^{\frac{4}{3}}}{z - x} \cdot \frac{z^{\frac{8}{3}} + z^{\frac{4}{3}}x^{\frac{4}{3}} + x^{\frac{8}{3}}}{z^{\frac{8}{3}} + z^{\frac{4}{3}}x^{\frac{4}{3}} + x^{\frac{8}{3}}}$$

$$= \lim_{z \rightarrow x} \frac{z^4 - x^4}{(z-x)(z^{\frac{8}{3}} + z^{\frac{4}{3}}x^{\frac{4}{3}} + x^{\frac{8}{3}})}$$

$$= \frac{4x^3}{3x^{\frac{8}{3}}} = \frac{4}{3}x^{\frac{1}{3}}$$

In fact, (*) holds for $n \in \mathbb{R}$
 (any real number)
 We will show it in next section.

Ex: $f(x) = x^4 - 2x^2 + 2$,
find all horizontal tangents
of $y = f(x)$.

Ans: Solve x from $f'(x) = 0$

$$f'(x) = 4x^3 - 4x + 0$$

(we have used the rule
 $\frac{d}{dx}(u(x) + v(x)) = \frac{du}{dx} + \frac{dv}{dx}$)

$$\Rightarrow x = 0, \pm 1$$

horizontal tangents at $(0, f(0))$
 $(1, f(1))$ and $(-1, f(-1))$ are

$$\frac{y-2}{x-0} = 0; \quad \frac{y-1}{x-1} = 0; \quad \frac{y-1}{x+1} = 0$$

Derivative Rules:

$$\frac{d}{dx}(C \cdot u(x)) = C \frac{d}{dx} u(x) \quad (1)$$

$C = \text{constant}$

$$\frac{d}{dx}(u(x) \pm v(x)) = \frac{du}{dx} \pm \frac{dv}{dx} \quad (2)$$

$$\frac{d}{dx}(u(x) \cdot v(x)) = u'v + uv' \quad (3)$$

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v u' - u v'}{v^2} \quad (4)$$

pf: (1): exercise

$$(2): (u+v)'$$

$$= \lim_{h \rightarrow 0} \frac{(u(x+h) + v(x+h)) - (u(x) + v(x))}{h}$$

$$= \lim_{h \rightarrow 0} \frac{u(x+h) - u(x)}{h} + \lim_{h \rightarrow 0} \frac{v(x+h) - v(x)}{h}$$

$$= u' + v' \quad (\text{similarly for (2)})$$

Differentiation Rules

pf of (3).

$$\begin{aligned}(uv)' &= \lim_{h \rightarrow 0} \frac{u(x+h)v(x+h) - u(x)v(x)}{h} \\&= \lim_{h \rightarrow 0} \left(\frac{u(x+h)v(x+h) - u(x+h)v(x)}{h} + \frac{u(x+h)v(x) - u(x)v(x)}{h} \right) \\&= \lim_{h \rightarrow 0} u(x+h) \left(\frac{v(x+h) - v(x)}{h} \right) \\&\quad + \lim_{h \rightarrow 0} v(x) \left(\frac{u(x+h) - u(x)}{h} \right) \\&= u(x)v'(x) + v(x)u'(x)\end{aligned}$$

pf of (4):

$$\frac{u}{v} = u \cdot \left(\frac{1}{v}\right)$$

It remains to evaluate

$$\left(\frac{1}{v}\right)' = \lim_{z \rightarrow x} \frac{\frac{1}{v(z)} - \frac{1}{v(x)}}{z - x}$$

$$= \lim_{z \rightarrow x} \frac{\left(\frac{v(x) - v(z)}{v(z)v(x)}\right)}{z - x}$$


$$= - \lim_{z \rightarrow x} \left(\frac{v(z) - v(x)}{z - x} \right) \frac{1}{v(z)v(x)}$$

$$= - \frac{v'}{v^2}$$

$$\begin{aligned} \text{From (3)} \Rightarrow \left(\frac{u}{v}\right)' &= u \left(\frac{1}{v}\right)' + u' \left(\frac{1}{v}\right) \\ &= \frac{uv' - uv'}{v^2} \end{aligned}$$

Exponential functions

$$y = a^x, \quad x \in \mathbb{R}, \quad a > 0$$



$$\frac{d}{dx} a^x = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{a^x \cdot a^h - a^x}{h}$$

$$= \left(\lim_{h \rightarrow 0} \frac{a^h - 1}{h} \right) \cdot a^x$$

$$= g(a) \cdot a^x$$


Here $g(a) = \lim_{h \rightarrow 0} \frac{a^h - 1}{h}$ (if exists)

$$g(1) = \lim_{h \rightarrow 0} \frac{1^h - 1}{h} = 0$$

$$\begin{aligned} g(a^2) &= \lim_{h \rightarrow 0} \frac{a^{2h} - 1}{h} \cdot \frac{2}{2} \\ &= \lim_{h \rightarrow 0} \frac{a^{2h} - 1}{2h} \cdot 2 = 2g(a) \end{aligned}$$

$a > 1 \Rightarrow g(a) \geq 0$ (In fact, it exists and > 0)

$a < 1 \Rightarrow g(a) \leq 0$ (< 0)



Summary:

$$(1) \frac{d}{dx} a^x = \left(\lim_{h \rightarrow 0} \frac{a^h - 1}{h} \right) \cdot a^x$$

$$(2) \text{ Let } e \approx 2.71828 \dots$$

$$\text{Satisfy } \left(\lim_{h \rightarrow 0} \frac{e^h - 1}{h} \right) = 1$$

$$\text{i.e. } \frac{d}{dx} e^x = e^x$$

Example $f(x) = e^{-x}$

$$f'(x) = ?$$

$$f(x) = \frac{1}{e^x}, \text{ from proof}$$

of (4) above

$$\begin{aligned} f'(x) &= \frac{-(e^x)'}{(e^x)^2} = \frac{-e^x}{e^{2x}} \\ &= -e^{-x} \end{aligned}$$